New Analytical and Numerical Results For Fractional Bogoyavlensky-Konopelchenko Equation Arising in Fluid Dynamics

Ali Kurt

Abstract. In this article, (2 + 1)-dimensional time fractional Bogoyavlensky-Konopelchenko (BK) equation is studied, which describes the interaction of wave propagating along the x axis and y axis. To acquire the exact solutions of BK equation we employed sub equation method that is predicated on Riccati equation, and for numerical solutions the residual power series method is implemented. Some graphical results that compares the numerical and analytical solutions are given for different values of μ . Also comparative table for the obtained solutions is presented.

§1 Introduction

Fractional calculus has been a desirable area of research for several years and is growing rapidly and ongoing development. As a branch of infinitesimal calculus, fractional calculus specially studies the possibility of performing fractional differentiation operator on functions, such as a derivative of order 1/2. In general, fractional calculus involves the following main advantages:

- 1. Fractional calculus can easily express the historical dependence of the evolution of system analysis by taking the global correlation into consideration, but integer calculus is not convenient to represent this process for its locality characteristics.
- 2. Fractional calculus overcomes the critical defect of integer calculus that the theoretical model results often fail to coincide with the experimental results. On the contrary, it could get a good coincidence by using a few parameters.
- 3. When describing complicated physical mechanics problems, fractional calculus has a clearer physical significance and a simpler expression compared with the nonlinear model.

Received: 2019-05-13. Revised: 2019-09-10.

MR Subject Classification: 35R11, 35A20, 35C05.

Keywords: Conformable Fractional Derivative, Fractional Bogoyavlensky-Konopelchenko Equation, Sub-Equation Method, Residual Power Series Method.

Digital Object Identifier(DOI): https://doi.org/10.1007/s11766-020-3808-9.

To benefit these advantages scientists expressed different definitions of fractional derivative and integrals. During the past four decades or so, various operators of fractional calculus are declared such as those named after Riemann-Liouville, Weyl, Hadamard, Grunwald-Letnikov, Riesz, Erdelyi-Kober, Liouville-Caputo, Caputo and so on [1-4]. Almost all of these operators involve the integral representation in their definition. This situation makes the calculations harder and the exact solution of the considered mathematical model can not be obtained. Also some of these definitions do not satisfy the basic properties to be a fractional derivative definition [5]. For instance the derivative of a arbitrary constant is not zero when the Riemann-Liouville is used as derivative operator. In addition Caputo and Riemann-Liouville definitions do not satisfy the rules for derivative of quotient of two functions, derivative of product of two functions, chain rule and etc. Recently a new fractional operator called "conformable fractional derivative and integral" is introduced by Khalil et al. [6] which overcomes the above mentioned deficiencies.

Definition Let $f:[0,\infty) \to \mathbb{R}$ be a function. The μ^{th} order "conformable fractional derivative" (CFD) [6] of f is defined as,

$$D_{\mu}(f)(t) = \lim_{\varepsilon \to 0} \frac{f(t + \varepsilon t^{1-\mu}) - f(t)}{\varepsilon},$$

for all $t > 0, \mu \in (0, 1)$. Let f be a μ -differentiable function in some (0, a), a > 0 and $\lim_{t \to 0^+} f^{(\mu)}(t)$ exists then $f^{(\mu)}(0) = \lim_{t \to 0^+} f^{(\mu)}(t)$. The "conformable fractional integral" of f starting from $a \ge 0$ is defined as:

$$I^a_\mu(f)(t) = \int\limits_a^t \frac{f(x)}{x^{1-\mu}} dx,$$

where the integral indicates the usual Riemann improper integral, and $\mu \in (0, 1]$. The CFD satisfies the properties that are given in the following theorem [6].

Theorem 1.1. Let $\mu \in (0,1]$ and suppose f, g are μ -differentiable at point t > 0. Then

- 1. $D_{\mu}(cf + dg) = cD_{\mu}(f) + cD_{\mu}(g)$ where Let $c, f \in \mathbb{R}$.
- 2. $D_{\mu}(t^p) = pt^{p-\mu}$ where $p \in \mathbb{R}$.
- 3. Let λ be a constant then $D_{\mu}(\lambda) = 0$.

4.
$$D_{\mu}(fg) = fD_{\mu}(g) + gD_{\mu}(f)$$

5.
$$D_{\mu}\left(\frac{f}{g}\right) = \frac{gD_{\mu}(f) - fD_{\mu}(g)}{g^2}$$

After this definition was expressed, which can be described as a milestone for fractional calculus, huge amount of articles has been done that involves conformable fractional derivative and integral [7–10,26,30,31]. For instance Javeed et al. [11] constructed the solution of conformable fractional Burger-Poisson equation by using homotopy perturbation method. Rosales et al. [12] employed conformable fractional derivative to analyze the classical Drude model. Alharbi et al. [28] investigated the projectile motion with the aid of conformable fractional derivative.

Yaslan and Girgin [14] implemented G'/G^2 expansion method to get the exact solutions of conformable time fractional Caudrey-Dodd-Gibbon equation, Calogero-Bogoyavlenskii-Schiff (CBS) equation and Ablowitz-Kaup-Newell-Segur equations. Cenesiz et al. [15] established the exact and numerical solutions of fractional Hirota-Satsuma Coupled KdV System where the fractional terms are in conformable sense. Also many researches including conformable derivative can be cited therein [27–29].

In this article the new analytical and numerical solutions of (2 + 1) dimensional time fractional BK Equation

$$D_t^{\mu} D_x u + \alpha D_x^4 u + \beta D_x^3 D_y u + 6\alpha D_x^2 u D_x u + 4\beta D_x D_y u D_x u + 4\beta D_x^2 u D_y u = 0, \qquad (1)$$

that identifies the interaction of a Riemann wave propagating along the y-axis and a long wave propagating along the x-axis in a fluid where u is a function of the space coordinates x and yand time coordinate t and the parameters α and β are constant. To the best of our knowledge all the obtained results are seen for the first time in the literature and will benefit the scientists that study on wave propagation.

§2 Brief Description of Implemented Methods

2.1 Sub Equation Method

Lets describe the considered method called sub equation method [16] which established on the Riccati equation

$$\varphi'(\xi) = \sigma + \left(\varphi(\xi)\right)^2. \tag{2}$$

The general form of nonlinear time fractional partial differential equation is

$$P\left(u, D_t^{\mu}u, D_x u, D_y u, D_x^2 u, D_y^2 u, D_x D_y u...\right) = 0,$$
(3)

where $D_t^{\mu} u$ denotes conformable derivative operator with fractional order. Regarding the fractional wave transformation [17]

$$u(x, y, t) = U(\xi), \ \xi = kx + ly + w \frac{t^{\mu}}{\mu},$$
(4)

where k, w, l are any constants to be evaluated later and the chain rule [18], Eq. (3) becomes an integer order nonlinear ordinary differential equation

$$G(U(\xi), U'(\xi), U''(\xi), ...) = 0.$$
(5)

Suppose that the solution of the reduced Eq. (5) in the following form

$$U(\xi) = \sum_{i=0}^{N} a_i \varphi^i(\xi), \ a_N \neq 0,$$
(6)

where a_i $(0 \le i \le N)$ are stable coefficients to be examined and N is can be acquired by balancing principle [19] in Eq. (5) and $\varphi(\xi)$ is the solution of Riccati Eq. (2). A set of the solutions which verify the Eq. (2) is given below.

Appl. Math. J. Chinese Univ. Vol. 35, No. 1

$$\varphi(\xi) = \begin{cases} -\sqrt{-\sigma} \tanh\left(\sqrt{-\sigma}\xi\right), \ \sigma < 0\\ -\sqrt{-\sigma} \coth\left(\sqrt{-\sigma}\xi\right), \ \sigma < 0\\ \sqrt{\sigma} \tan\left(\sqrt{\sigma}\xi\right), \ \sigma > 0\\ -\sqrt{\sigma} \cot\left(\sqrt{\sigma}\xi\right), \ \sigma > 0\\ -\frac{1}{\xi+\varpi}, \ \varpi \ is \ a \ cons., \ \sigma = 0 \end{cases}$$
(7)

Collecting the all the obtained data sets we attain a polynomial including $\varphi(\xi)$. Vanishing the coefficients of $\varphi^i(\xi)$ (i = 0, 1, ..., N) produces an algebraic system in k, l, w, a_i (i = 0, 1, ..., N). Calculating the solution of these nonlinear algebraic equation we obtain the values for l, w, k, a_i (i = 0, 1, ..., N). Substituting all the obtained values in the formulas (7) we get the exact solutions for Eq. (3).

2.2 Residual Power Series Method

To illustrate the idea of RPSM [20–23], one can regard the following equation [24]:

$$T_{\mu}u(x,y,t) + N[x,y]u(x,y,t) + L[x,y]u(x,y,t) = c(x,y,t),$$
(8)

where $x \in \mathbb{R}, n - 1 < n\mu \leq n, t > 0$ and given with the initial condition

$$f_0(x,y) = u(x,y,0) = f(x,y).$$
(9)

where the operator L[x, y] is linear, the operator N[x, y] is nonlinear and c(x, y, t) are arbitrary function.

The RPSM established on the indicating the solution of the Eq. (8) according to (9) with fractional power series expansion around t = 0.

$$f_{(n-1)}(x,y) = T_t^{(n-1)\mu} u(x,y,0) = h(x,y).$$
(10)

The expanded version of the solution can be regarded as

$$u(x,y,t) = f(x,y) + \sum_{n=1}^{\infty} f_n(x,y) \frac{t^{n\mu}}{\mu^n n!}.$$
(11)

Then, $u_k(x, y, t)$ can be written as:

$$u_k(x, y, t) = f(x, y) + \sum_{n=1}^k f_n(x, y) \frac{t^{n\mu}}{\mu^n n!}.$$
(12)

Since the 1st approximate solution $u_1(x, y, t)$ is

$$u_1(x, y, t) = f(x, y) + f_1(x, y) \frac{t^{\mu}}{\mu},$$
(13)

so $u_k(x, y, t)$ can be rearranged as follows

$$u_k(x,y,t) = f(x,y) + f_1(x,y)\frac{t^{\mu}}{\mu} + \sum_{n=2}^k f_n(x,y)\frac{t^{n\mu}}{\mu^n n!}, \ k = 2,3,4,\dots$$
(14)

for $0 < \mu \leq 1, 0 \leq t < \mathbb{R}^{\frac{1}{v}}, x \in I$.

Initially we express the residual function and the k - th residual function

104

Ali Kurt.

$$Res(x, y, t) = T_{\mu}u(x, y, t) + N[x, y]u(x, y, t) + L[x, y]u(x, y, t) - c(x, y, t),$$
(15)

 $Res_k(x,t) = T_{\mu}u_k(x,y,t) + N[x,y]u_k(x,y,t) + L[x,y]u_k(x,y,t) - c(x,y,t), \ k = 1,2,3,\dots \ (16)$ respectively. Obviously, Res(x, y, t) = 0 and $\lim_{k \to \infty} Res_k(x, y, t) = Res(x, y, t)$ for each $x \in I$ and $t \ge 0$. Indeed this bring about $\frac{\partial^{(n-1)\mu}}{\partial t^{(n-1)\mu}} Res_k(x, y, t) = 0$ for n = 1, 2, 3, ..., k [22, 23, 25]. Evaluating the solution of the equation $\frac{\partial^{(n-1)\mu}}{\partial t^{(n-1)\mu}} Res_k(x, y, 0) = 0$ concludes the necessary $f_n(x,y)$ coefficients. Thus the numerical solutions can be acquired respectively.

Analytical Solutions of the Bogoyavlensky-Konopelchenko §3 Equation

Consider the time fractional BK equation

$$D_t^{\mu} D_x u + \alpha D_x^4 u + \beta D_x^3 D_y u + 6\alpha D_x^2 u D_x u + 4\beta D_x D_y u D_x u + 4\beta D_x^2 u D_y u = 0, \quad (17)$$

where $D_t^{\mu} u$ is conformable derivative of function $u(x, y, t)$ with fractional order. Implementing
the chain [18] and wave transform (4) to Eq. (17), the reduced form can be expressed as follows

$$kwU' + (\alpha k^4 + \beta k^3 l)U''' + (3\alpha k^3 + 4\beta k^2 l)(U')^2 = 0.$$
 (18)

Regard that the solution of Eq. (18) is in terms of $\varphi(\xi)$ where is the exact solutions of Eq. (2) as follows.

$$U(\xi) = \sum_{i=0}^{N} a_i \varphi^i(\xi), \ a_N \neq 0.$$
 (19)

With use of the balancing principle [19], we have N = 1. Gathering all the obtained data in Eq. (18), an algebraic equation system come to exist with respect to k, l, w, a_0, a_1 . Solving the obtain system yields following solution set

$$a_1 = \frac{3kw}{2\left(\alpha k^3 \sigma - w\right)}, l = \frac{w - 4\alpha k^3 \sigma}{4\beta k^2 \sigma},\tag{20}$$

and k, w, a_0 are free constants. When $\sigma < 0$, using (7) and (4) the traveling wave solutions of Eq. (17) can be deducted

$$u_1(x, y, t) = a_0 - \frac{3k\sqrt{-\sigma}w\tanh\left(\sqrt{-\sigma}\xi\right)}{2\left(\alpha k^3\sigma - w\right)},$$
$$u_2(x, y, t) = a_0 - \frac{3k\sqrt{-\sigma}w\coth\left(\sqrt{-\sigma}\xi\right)}{2\left(\alpha k^3\sigma - w\right)},$$

where $\xi = kx + \frac{(w - 4\alpha k^3 \sigma)}{4\beta k^2 \sigma}y + \frac{wt^{\mu}}{\mu}$. In a similar way, for $\sigma > 0$, we obtain

$$\begin{split} u_3(x,y,t) &= a_0 + \frac{3k\sqrt{\sigma}w\tan\left(\sqrt{\sigma}\xi\right)}{2\left(\alpha k^3\sigma - w\right)},\\ u_4(x,y,t) &= a_0 - \frac{3k\sqrt{\sigma}w\cot\left(\sqrt{\sigma}\xi\right)}{2\left(\alpha k^3\sigma - w\right)},\\ \end{split}$$
 here $\xi &= kx + \frac{\left(w - 4\alpha k^3\sigma\right)}{4\beta k^2\sigma}y + \frac{wt^{\mu}}{\mu}. \end{split}$

wł

§4 Approximate Solutions of the Bogoyavlensky-Konopelchenko Equation

Consider the nonlinear time-fractional BK Eq. (17). The initial condition obtained from the exact solution is

$$u(x, y, 0) = a_0 - \frac{3k\sqrt{-\sigma}w\tanh\left(\sqrt{-\sigma}\left(\frac{y(w-4\alpha k^3\sigma)}{4\beta k^2\sigma} + kx\right)\right)}{2\left(\alpha k^3\sigma - w\right)}.$$

For residual power series

$$u(x, y, t) = f(x, y) + \sum_{n=1}^{\infty} f_n(x, y) \frac{t^{n\mu}}{\mu^n n!},$$

and k.truncated series of u(x,t)

$$u_k(x, y, t) = f(x, y) + \sum_{n=1}^k f_n(x, y) \frac{t^{n\mu}}{\mu^n n!}, \ k = 1, 2, 3, ..$$

Therefore, the k-th residual functions of time-fractional B-K equation is: $\begin{aligned} Resu_k(x, y, t) &= (\partial_t^{\mu} u_k)_x + \alpha \, (u_k)_{xxxx} + \beta \, (u_k)_{xxxy} + 6\alpha \, (u_k)_{xx} \, (u_k)_x + 4\beta \, (u_k)_{xx} \, (u_k)_y \\ &+ 4\beta \, (u_k)_{xy} \, (u_k)_x \,. \end{aligned}$

To determine the coefficients $f_1(x, y)$, in $u_1(x, y, t)$, we should replace the 1st truncated series $u_1(x, y, t) = f(x, y) + f_1(x, y) \frac{t^{\mu}}{\mu}$ into the 1st truncated residual function as

$$\begin{aligned} Resu_{1}(x, y, t) &= 6\alpha \left(f_{x} + \frac{t^{\mu} (f_{1})_{x}}{\mu} \right) \left(f_{xx} + \frac{t^{\mu} (f_{1})_{xx}}{\mu} \right) \\ &+ \alpha \left(f_{xxxx} + \frac{t^{\mu} (f_{1})_{xxxx}}{\mu} \right) \\ &+ 4\beta \left(f_{x}(x, y) + \frac{t^{\mu} (f_{1})_{x}}{\mu} \right) \left(f_{xy}(x, y) + \frac{t^{\mu} (f_{1})_{xy}}{\mu} \right) \\ &+ 4\beta \left(f_{y} + \frac{t^{\mu} (f_{1})_{y}}{\mu} \right) \left(f_{xx} + \frac{t^{\mu} (f_{1})_{xx}}{\mu} \right) \\ &+ \beta \left(f_{xxxy} + \frac{t^{\mu} (f_{1})_{xxxy}}{\mu} \right) + (f_{1})_{x} \,, \end{aligned}$$

for f = f(x, y). Now for the substitution of t = 0 through the equation $Resu_1(x, y, t)$ to obtain

$$(f_1)_x = -6\alpha f_x f_{xx} - \alpha f_{xxxx} - 4\beta f_x f_{xy} - 4\beta f_y f_{xx} - \beta f_{xxxy}.$$

Solving this differential equation gives the first unknown parameter as

$$f_1 = -3\alpha f_x^2 - \alpha f - 4\beta f_x f_y - \beta f_{xxy}.$$

Thus, we obtain the 1st RPS approximate solutions of time-fractional B-K equation as

$$u_1(x,y,t) = f + \frac{t^{\mu} \left(-3\alpha f_x^2 - \alpha f_{xxx} - 4\beta f_x f_y - \beta f_{xxy}\right)}{\mu}.$$

Again, to determine the second unknown coefficient $f_2(x, y)$, we replace the 2nd truncated series solution $u_2(x, y, t) = f(x, y) + f_1(x, y) \frac{t^{\mu}}{\mu} + f_2(x, y) \frac{t^{2\mu}}{2\mu^2}$ into the 2nd truncated residual

function and obtain

$$\begin{aligned} Resu_{2}(x,y,t) &= 6\alpha \left(f_{x} + \frac{t^{\mu} (f_{1})_{x}}{\mu} + \frac{t^{2\mu} (f_{2})_{x}}{2\mu^{2}} \right) \left(f_{xx} + \frac{t^{2\mu} (f_{2})_{xx}}{2\mu^{2}} + \frac{t^{\mu} (f_{1})_{xx}}{\mu} \right) \\ &+ \alpha \left(f_{xxxx} + \frac{t^{\mu} (f_{1})_{xxxxx}}{\mu} + \frac{t^{2\mu} (f_{2})_{xxxxx}}{2\mu^{2}} \right) \\ &+ 4\beta \left(f_{x} + \frac{t^{\mu} (f_{1})_{x}}{\mu} + \frac{t^{2\mu} (f_{2})_{x}}{2\mu^{2}} \right) \\ &\times \left(f_{xy} + \frac{t^{\mu} (f_{1})_{xy}}{\mu} + \frac{t^{2\mu} (f_{2})_{xy}}{2\mu^{2}} \right) + 4\beta \left(f_{y} + \frac{t^{\mu} (f_{1})_{y}}{\mu} + \frac{t^{2\mu} (f_{2})_{y}}{2\mu^{2}} \right) \\ &\times \left(f_{xx} + \frac{t^{\mu} (f_{1})_{xxx}}{\mu} + \frac{t^{2\mu} (f_{2})_{xxxy}}{2\mu^{2}} \right) \\ &+ \beta \left(f_{xxxy} + \frac{t^{\mu} (f_{1})_{xxxy}}{\mu} + \frac{t^{2\mu} (f_{2})_{xxxy}}{2\mu^{2}} \right) \\ &+ \frac{t^{\mu} (f_{2})_{x}}{\mu} + (f_{1})_{x}. \end{aligned}$$

Now, applying T_{μ} on both sides of $Resu_2(x, y, t)$ and equating to 0 for t = 0 gives:

$$(f_2)_x = -6\alpha (f_1)_x f_{xx} - 6\alpha f_x (f_1)_{xx} - 4\beta (f_1)_x f_{xy} - 4\beta f_x \times (f_1)_{xy} - 4\beta (f_1)_y f_{xx} - 4\beta f_y (f_1)_{xx} - \alpha (f_1)_{xxxx} - \beta (f_1)_{xxxy}.$$

Solving this differential equation gives

$$f_2(x,y) = -2 (f_1)_{xy} (3\alpha f_x + 2\beta f_y) - 4\beta (f_1)_x (f_1)_y - \alpha (f_1)_{xxx} - \beta (f_1)_{xxy}.$$

The 2nd RPS approximate solutions of time-fractional B-K equation is:

$$u_{2}(x,y,t) = f + \frac{t^{\mu}f_{1}}{\mu} + \frac{t^{2\mu}\left(-2\left(f_{1}\right)_{x}\left(3\alpha f_{x} + 2\beta f_{y}\right) - 4\beta f_{x}\left(f_{1}\right)_{y} - \alpha\left(f_{1}\right)_{xxx} - \beta\left(f_{1}\right)_{xxy}\right)}{2\mu^{2}}.$$

In a similar way, operating the same calculation for n = 3 and 4 to get the following results.

$$\begin{split} f_{3}(x,y) &= -2 \left(f_{2}\right)_{x} \left(3\alpha f_{y} + 2\beta f_{y}\right) - 4\beta \left(f_{2}\right)_{y} f_{x} - 6\alpha \left(f_{1}^{2}\right)_{x} \\ &-\alpha \left(f_{2}\right)_{xxx} - 8\beta \left(f_{1}\right)_{x} \left(f_{1}\right)_{y} - \beta \left(f_{2}\right)_{xxy}\right), \\ u_{3}(x,y,t) &= f + \frac{t^{\mu} f_{1}}{\mu} + \frac{t^{2\mu} f_{2}}{2\mu^{2}} \\ &+ \frac{t^{3\mu} \left(-2 \left(f_{2}\right)_{x} \left(3\alpha f_{x} + 2\beta f_{y}\right) - 4\beta \left(f_{2}\right)_{y} f_{x} - 6\alpha \left(f_{1}^{2}\right)_{x}\right)}{6\mu^{3}} \\ &+ \frac{t^{3\mu} \left(-\alpha \left(f_{2}\right)_{xxx} - 8\beta \left(f_{1}\right)_{x} \left(f_{1}\right)_{y} - \beta \left(f_{2}\right)_{xxy}\right)}{6\mu^{3}}, \\ f_{4}(x,y) &= -2 \left(f_{3}\right)_{x} \left(3\alpha f_{x} + 2\beta f_{y}\right) - 4\beta \left(f_{3}\right)_{y} f_{x} - 6 \left(f_{2}\right)_{x} \left(3\alpha \left(f_{1}\right)_{x} + 2\beta \left(f_{1}\right)_{y}\right) - \alpha \left(f_{3}\right)_{xxx} \end{split}$$

Appl. Math. J. Chinese Univ.

Vol. 35, No. 1

$$u_{4}(x,y,t) = f + \frac{t^{\mu}f_{1}}{\mu} + \frac{t^{2\mu}f_{2}}{2\mu^{2}} + \frac{t^{3\mu}f_{3}}{6\mu^{3}} + \frac{t^{4\mu}}{24\mu^{4}} \left(-2(f_{3})_{x}(3\alpha f_{x} + 2\beta f_{y}) - 4\beta(f_{3})_{y}f_{x}\right) - 6(f_{2})_{x}\left(3\alpha(f_{1})_{x} + 2\beta(f_{1})_{y}\right) + \frac{t^{4\mu}\left(-\alpha(f_{3})_{xxx} - 12\beta(f_{2})_{y}(f_{1})_{x} - \beta(f_{3})_{xxy}\right)}{24\mu^{4}}.$$

In Table 1, the fourth-order approximate RPSM solutions of time-fractional B-K equation are compared numerically with the exact solution

$$u(x,y,t) = a0 - \frac{3k\sqrt{-\sigma}w\tanh\left(\sqrt{-\sigma}\left(\frac{y(w-4\alpha k^{3}\sigma)}{4\beta k^{2}\sigma} + kx + \frac{wt^{\mu}}{\mu}\right)\right)}{2\left(\alpha k^{3}\sigma - w\right)}$$

The graphical representations indicates that obtained exact solutions and numerical results are compatible for different values of μ . Also the table that given below shows that the absolute error values are in acceptable limits.

Table 1: RPSM approximate results and comparison with the exact solutions by absolute errors for $a_0 = 1$, k = 2, w = 3, $\sigma = -4$, $\alpha = 2.5$, $\beta = 1$, and t = 0.1.

	1								
	$\mu = 0.50$			$\mu = 0.75$			$\mu = 0.95$		
x	RPSM	Exact	Abs. Error	RPSM	Exact	Abs. Error	RPSM	Exact	$Abs. \ Error$
0.0	0.783133	0.783134	1.28127E-6	0.783133	0.783133	2.03316E-9	0.783133	0.783133	4.5705E-11
0.1	0.783133	0.783136	2.85150E-6	0.783133	0.783133	4.52488E-9	0.783133	0.783133	1.0171E-10
0.2	0.783133	0.783140	6.34607E-6	0.783133	0.783133	1.00703E-8	0.783133	0.783133	2.2638E-10
0.3	0.783135	0.783149	1.41231E-5	0.783133	0.783133	2.24119E-8	0.783133	0.783133	5.0381E-10
0.4	0.783137	0.783168	3.14299E-5	0.783133	0.783133	4.98785E-8	0.783133	0.783133	1.12127E-9
0.5	0.783143	0.783213	6.99404E-5	0.783133	0.783133	1.11006E-7	0.783133	0.783133	2.49542E-9
0.6	0.783155	0.783311	1.55615E-4	0.783134	0.783134	2.47047E-7	0.783133	0.783133	5.55363E-9
0.7	0.783182	0.783528	3.46129E-4	0.783135	0.783136	5.49802E-7	0.783133	0.783133	1.23597E-8
0.8	0.783243	0.784013	7.69339E-4	0.783139	0.783140	1.22356E-6	0.783134	0.783134	2.75063E-8
0.9	0.783379	0.785087	1.70734E-3	0.783147	0.783150	2.72283E-6	0.783137	0.783137	6.12131E-8
1.0	0.783681	0.787457	3.77590E-3	0.783164	0.783171	6.05853E-6	0.783142	0.783142	1.36216E-7



Figure 1: RPSM solution and exact solution for $a_0 = 1$, k = 2, w = 3, $\sigma = -4$, $\alpha = 2.5$, $\beta = 1$, t = 0 and $\mu = 0.50$.

108



Figure 2: RPSM solution and exact solution for $a_0 = 1$, k = 2, w = 3, $\sigma = -4$, $\alpha = 2.5$, $\beta = 1$, t = 0 and $\mu = 0.75$.



Figure 3: RPSM solution and exact solution for $a_0 = 1$, k = 2, w = 3, $\sigma = -4$, $\alpha = 2.5$, $\beta = 1$, t = 0 and $\mu = 0.95$.

§5 Conclusion

In this article the author obtained the numerical and analytical solutions of conformable time fractional BK equation that arises in interaction of the wave propagation with the aid of sub equation method and RPSM. The 3D graphical representation for comparisons of the obtained results are given for different values of μ . Also comparative table is represented to express the effectiveness, reliableness and accuracy of the implemented methods. All the exact, numerical solutions and also, graphical representations, tables are obtained with the aid of computer software called Mathematica. This work reported here is a first step towards understanding structural and physical behaviour of the models arising in fluid dynamics. We hope that this work will be very useful in better understanding the physical structure occurring at real life events that correspond to the models of fluid dynamics. We believe our manuscript is very timely and will interest a broad range of scientists that work on mathematical models arising in fluid dynamics.

References

- K S Miller, B Ross. An Introduction to the Fractional Calculus and Fractional Differential Equations, John Wiley & Sons, New York, 1993.
- [2] A Kilbas, H M Srivastava, J J Trujillo. Theory and Applications of Fractional Differential Equations, Elsevier, San Diego, 2006.
- [3] I Podlubny. Fractional Differential Equations, Academic Press, San Diego, 1999.
- [4] S Das. Functional fractional calculus, Springer, 2011.
- [5] A Atangana. Derivative with a new parameter: Theory, methods and applications, Academic Press, 2015.
- [6] R Khalil, M Al Horani, A Yousef, M Sababheh. A new definition of fractional derivative, J Comput Appl Math, 2017, 264: 65-70.
- [7] A Kurt, O Tasbozan, D Baleanu. New solutions for conformable fractional NizhnikNovikovVeselov system via G'/G expansion method and homotopy analysis methods, Opt Quant Electron, 2017, 49: 333.
- [8] O Tasbozan, M Senol, A Kurt, O Ozkan. New solutions of fractional Drinfeld-Sokolov-Wilson system in shallow water waves, Ocean Eng, 2018, 161: 62-68.
- [9] O Tasbozan, Y Cenesiz, A Kurt, D Baleanu. New analytical solutions for conformable fractional PDEs arising in mathematical physics by exp-function method, Open Phys, 2017, 15: 647-651.

- [10] A Kurt, O Tasbozan, Y Cenesiz. Homotopy analysis method for conformable BurgersKorteweg-de Vries equation, Bull Math Sci Appl, 2016, 17: 17-23.
- [11] S Javeed, D Baleanu, A Waheed, M Shaukat Khan, H Affan. Analysis of Homotopy Perturbation Method for Solving Fractional Order Differential Equations, Mathematics, 2019, 7(1): 40.
- [12] J J Rosales, F A Godnez, V Banda, G H Valencia. Analysis of the Drude model in view of the conformable derivative, Optik, 2019, 178: 1010-1015.
- [13] F M Alharbi, D Baleanu, A Ebaid. Physical properties of the projectile motion using the conformable derivative, Chinese J Phys, 2019, 58: 18-28.
- [14] H C Yaslan, A Girgin. New exact solutions for the conformable space-time fractional KdV, CDG, (2+1)-dimensional CBS and (2+1)-dimensional AKNS equations, J Taibah Univ Sci, 2019, 13: 1-8.
- [15] Y Cenesiz, A Kurt, O Tasbozan. On the New Solutions of the Conformable Time Fractional Generalized Hirota-Satsuma Coupled KdV System, Ann West Univ Timisoara-Math Comput Sci, 2017, 55: 37-50.
- [16] S Zhang, H Q Zhang. Fractional sub-equation method and its applications to nonlinear fractional PDEs, Phys Lett A, 2011, 375: 1069-1073.
- [17] Y Cenesiz, A Kurt. New fractional complex transform for conformable fractional partial differential equations, J Appl Mat Stat Inform, 2016, 12: 41-47.
- [18] T Abdeljawad. On conformable fractional calulus, J Comput Appl Math, 2015, 279: 57-66.
- [19] W Malfliet. Solitary wave solutions of nonlinear wave equations, Am J Phys, 1992, 60: 650-654.
- [20] M Alquran. Analytical solutions of fractional foam drainage equation by residual power series method, Math Sci, 2015, 8: 153.
- [21] M Alquran. Analytical solution of time-fractional two-component evolutionary system of order 2 by residual power series method, J Appl Anal Comput, 2015, 5: 89-599.
- [22] O A Arqub. Series solution of fuzzy differential equations under strongly generalized differentiability, J Adv Res Appl Math, 2013, 5: 31-52.
- [23] O A Arqub, A El-Ajou, A Bataineh, I Hashim. A representation of the exact solution of generalized Lane Emden equations using a new analytical method, Abstr Appl Anal, 2013, Article ID 378593, 10 pages.
- [24] A Kumar, S Kumar, M Singh. Residual power series method for fractional Sharma-Tasso-Olever equation, Commun Numer Anal, 2016, 1: 1-10.

- [25] H M Jaradat, S Al-Shara, Q J Khan, M Alquran, K Al-Khaled. Analytical solution of timefractional Drinfeld-Sokolov-Wilson system using residual power series method, IAENG Int J Appl Math, 2016, 46: 64-70.
- [26] H C Yaslan. Numerical solution of the conformable space-time fractional wave equation, Chinese J Phys, 2018, 56(6): 2916-2925.
- [27] H Rezazadeh, H Tariq, M Eslami, M Mirzazadeh, Q Zhou. New exact solutions of nonlinear conformable time-fractional Phi-4 equation, Chinese J Phys, 2018, 56(6): 2805-2816.
- [28] F M Alharbi, D Baleanu, A Ebaid. Physical properties of the projectile motion using the conformable derivative, Chinese J Phys, 2019, 58: 18-28.
- [29] D Kumar, A R Seadawy, A K Joardar. Modified Kudryashov method via new exact solutions for some conformable fractional differential equations arising in mathematical biology, Chinese J Phys, 2018, 56(1): 75-85.
- [30] Javeed, S Riaz, K Saleem Alimgeer, M Atif, A Hanif, D Baleanu. First Integral Technique for Finding Exact Solutions of Higher Dimensional Mathematical Physics Models, Symmetry, 2019, 11(6): 783.
- [31] S Javeed, S Saif, A Waheed, D Baleanu. Exact solutions of fractional mBBM equation and coupled system of fractional Boussinesq-Burgers, Results Phys, 2018, 9: 275-1281.
- Department of Mathematics, Pamukkale University, Denizli 20180, Turkey. Email: pau.dr.alikurt@gmail.com, akurt@pau.edu.tr