

Spreading speed of a food-limited population model with delay

TIAN Ge^{1,2,*} AN Ruo-fan³

Abstract. This paper is concerned with the spreading speed of a food-limited population model with delay. First, the existence of the solution of Cauchy problem is proved. Then, the spreading speed of solutions with compactly supported initial data is investigated by using the general Harnack inequality. Finally, we present some numerical simulations and investigate the dynamical behavior of the solution.

§1 Introduction

In this paper, we consider the following food-limited population model with delay:

$$v_t(t, x) = \Delta v(t, x) + \mu v(t, x) \left(\frac{1 - v(t - \tau, x)}{1 + \beta v(t - \tau, x)} \right), \quad t \geq 0, \quad x \in \mathbb{R}, \quad (1.1)$$

where the delay τ and the coefficients μ, β are positive. The archetype of this model can be found in the original paper of Smith [18]. We also refer the readers to the more detailed research about the food-limited model in [3,4,6-10,13,17,19]. In particular, there are some works [8,12,22,25,26,28] about the traveling wave solution of (1.1). It is worth noting that Trofimchuk et al. [22] presented a broad and explicit range of the parameters which permit the existence of monotone traveling waves. Furthermore, Hasik et al. [12] gave a more detailed analysis about the geometric diversity of wavefront of (1.1), which include the non-monotone and non-oscillating wavefronts.

However, there seem no results about the long-time behaviour of solutions of the Cauchy problem of (1.1). The long-time behaviors of solutions of reaction-diffusion equations with compactly supported initial data can well describe the propagation phenomenon in population invasion and epidemic spread. Usually, such a propagation phenomenon can be characterized

Received: 2020-08-08. Revised: 2022-01-09.

MR Subject Classification: 35K57, 35B40, 35B51, 35R09, 92D25.

Keywords: food-limited population model, reaction-diffusion equations, delay, spreading speed.

Digital Object Identifier(DOI): <https://doi.org/10.1007/s11766-023-4232-8>.

Supported by the National Natural Science Foundation of China (11371179).

*Corresponding author.

by investigating the so-called spreading speed of solutions [1, §4]. For convenience, we first introduce the following definition:

Definition 1.1 (Spreading speed). *Assume that $v(t, x)$ is a nonnegative function for all $(t, x) \in (0, +\infty) \times \mathbb{R}$. Then, there exists a constant $\kappa > 0$, c^* is called the asymptotic speed of spreading of $v(t, x)$ if*

- (i) $\lim_{t \rightarrow +\infty, |x| \geq (c^* + \varepsilon)t} v(t, x) = 0$ for any given $\varepsilon > 0$;
- (ii) $\liminf_{t \rightarrow +\infty, |x| \leq (c^* - \varepsilon)t} v(t, x) > \kappa$ for any given $\varepsilon \in (0, c^*)$.

Up to now, there has been great progress on the study of the spreading speed of solutions with compactly supported initial data when the solution semiflow of the equations is monotone. Unfortunately, equation (1.1) is not quasi-monotone. Therefore, the whole point of this paper is to investigate the spreading speed of solutions of (1.1) with compactly supported initial data, namely, we consider the following initial valued problem:

$$\begin{cases} v_t(t, x) = \Delta v(t, x) + \mu v(t, x) \left(\frac{1 - v(t - \tau, x)}{1 + \beta v(t - \tau, x)} \right), & t \geq 0, x \in \mathbb{R}, \\ v(s, x) = v_0(s, x), & s \in [-\tau, 0], x \in \mathbb{R}. \end{cases} \quad (1.2)$$

Note that the delay in the reaction term makes the comparison principle unavailable, which results in that the abstract theory of the spreading speed for monotone semiflows is not applicable. Recently, replacing the delayed term $v(t - \tau, x)$ by a special convolution term

$$(g * v)(t, x) = \int_0^\tau k(s) \int_{\mathbb{R}} \frac{1}{\sqrt{4\pi s}} e^{-(x-y)^2/(4s)} v(t-s, y) dy ds,$$

we [23] have determined the spreading speed by using the solution of the heat equation. Unfortunately, the method used in [23] is not applicable to equation (1.2). In particular, when $\beta = 0$, equation (1.2) is reduced to the following delayed Fisher-KPP equation:

$$\begin{cases} v_t(t, x) = \Delta v(t, x) + \mu v(t, x) (1 - v(t - \tau, x)), & t \geq 0, x \in \mathbb{R}, \\ v(s, x) = v_0(s, x), & s \in [-\tau, 0], x \in \mathbb{R}. \end{cases} \quad (1.3)$$

It should be emphasized that determining of the spreading speed of solutions of (1.3) is still an open problem. Of course, if an instantaneous self-limitation term is introduced to the nonlinearity, the boundedness of the solutions could be obtained and hence, the spreading speed of solutions could be established [14,15].

In this paper, we develop a new argument to establish the spreading speed of (1.2), which gets rid of the intricate investigations about the uniform boundedness of the solutions of (1.2). Precisely speaking, we will use the general Harnack inequality [5] and the comparison principle to study the spreading speed of solutions of (1.2). Before stating the main results, we give some notations. Let $\mathcal{Y} = BUC(\mathbb{R}, \mathbb{R})$ be the Banach space of all bounded and uniformly continuous functions from \mathbb{R} to \mathbb{R} with the supremum norm $\|\cdot\|_{\mathcal{Y}}$. Let $\mathcal{Y}^+ := \{v \in \mathcal{Y} : v(x) \geq 0, x \in \mathbb{R}\}$. Then \mathcal{Y} is a Banach lattice under the partial ordering induced by \mathcal{Y}^+ . It follows from [2, Theorem

1.5] that the \mathcal{Y} -realization $\Delta_{\mathcal{Y}}$ of the Laplacian Δ generates a strongly continuous analytic semigroup $\Psi(t)$ on \mathcal{Y} and $\Psi(t)\mathcal{Y}^+ \subset \mathcal{Y}^+$ for $t \geq 0$. In addition, we have

$$(\Psi(t)w)(x) = \frac{1}{\sqrt{4\pi t}} \int_{\mathbb{R}} e^{-\frac{(x-y)^2}{4t}} w(y) dy, \quad t > 0, x \in \mathbb{R}, w(\cdot) \in \mathcal{Y}.$$

Let $\mathcal{C} = C([- \tau, 0], \mathcal{Y})$ be the Banach space of continuous functions from $[- \tau, 0]$ into \mathcal{Y} with the supremum norm $\|\cdot\|_{\mathcal{C}}$ and let $\mathcal{C}^+ = \{\varphi \in \mathcal{C} : \varphi(s) \in \mathcal{Y}^+, s \in [- \tau, 0]\}$. Then \mathcal{C}^+ is a positive cone of \mathcal{C} . Usually, we identify an element $\varphi \in \mathcal{C}$ as a function from $[- \tau, 0] \times \mathbb{R}$ into \mathbb{R} defined by $\varphi(s, x) = \varphi(s)(x)$. For $T > 0$, we define $\mathcal{C}_T = C([- \tau, T], \mathcal{Y})$ be the Banach space of continuous functions from $[- \tau, T]$ into \mathcal{Y} with the supremum norm $\|\cdot\|_{\mathcal{C}_T}$.

Theorem 1.2. *Let $v(t, x)$ be the solution of the Cauchy problem (1.2) with the initial condition $v_0 \in \mathcal{C}^+$ such that $v_0(s, x) \not\equiv 0$. Then there exists γ which is a positive constant independent of $v_0 \in \mathcal{C}^+$, such that*

$$\liminf_{t \rightarrow +\infty} \left(\min_{|x| \leq ct} v(t, x) \right) > \gamma^{-1}, \quad \forall 0 \leq c < 2\sqrt{\mu}. \quad (1.4)$$

Furthermore, if $v_0(s, x)$ is compactly supported, then

$$\lim_{t \rightarrow +\infty} \left(\max_{|x| \geq ct} v(t, x) \right) = 0, \quad \forall c > 2\sqrt{\mu}. \quad (1.5)$$

This paper is organized as follows: in Section 2, we give the rigorous proof of Theorem 1.2. In Section 3, some numerical simulations are presented to investigate the dynamical behavior of the solution of (1.2).

§2 Main proof

The aim of this section is to provide a mathematically rigorous proof of Theorem 1.2. First of all, we give the global existence of the solution of (1.4) by the contraction mapping principle, as well as the classical argument.

Proposition 2.1 (Existence). *For any $v_0 \in \mathcal{C}^+$, equation (1.2) admits a unique mild solution $v(t, x) = v(t)(x) \in C([- \tau, \infty), \mathcal{Y})$ satisfying*

$$0 \leq v(t, x) \leq e^{\mu t} \|v_0\|_{\mathcal{C}_0}, \quad \forall t > 0, x \in \mathbb{R}.$$

In particular, when $t > \tau$, the mild solution is classical.

Proof. We first prove that the solution of (1.2) exists locally. Let $v_0 \in C([- \tau, 0], \mathcal{Y}^+)$ be the given initial value. For $0 < T < \infty$, fix $K > 1$, then we define

$$\Gamma_T := \left\{ v(t, x) \in \mathcal{C}_T : \begin{array}{l} v(s, x) = v_0(s, x) \text{ for } (s, x) \in [- \tau, 0] \times \mathbb{R}; \\ 0 \leq v(t, x) \leq K \|v_0\|_{\mathcal{C}_0} \text{ for } (t, x) \in [0, T] \times \mathbb{R}. \end{array} \right\}$$

It is clear that Γ_T is a complete metric space with a distance induced by the norm $\|\cdot\|_{\mathcal{C}_T}$. Let $L > \frac{\mu}{\beta}$ be a constant. For any $v \in \Gamma_T$, define

$$G(v)(t, x) := v(t, x) \left(L + \mu \frac{(1 - v(t - \tau, x))}{1 + \beta v(t - \tau, x)} \right)$$

and

$$u(t, x) := e^{-Lt}(\Psi(t)v_0(\cdot))(x) + \int_0^t e^{-L(t-s)}(\Psi(t-s)G(v)(s, \cdot))(x)ds \quad (2.1)$$

for $t \in [0, T]$ and $x \in \mathbb{R}$. According to [2,16,24], the function $u \in C([0, T], \mathcal{Y})$ defined by **representation** (2.1) is a mild solution of the following Cauchy problem

$$\begin{cases} u_t(t, x) - \Delta u(t, x) + Lu(t, x) = G(v)(t, x), & \forall t \in (0, T], x \in \mathbb{R}, \\ u(0, x) = v_0(x), & \forall x \in \mathbb{R}. \end{cases} \quad (2.2)$$

Define a map $\mathcal{T} : \Gamma_T \rightarrow \mathcal{C}_T$ by $\hat{u}(t, x) := (\mathcal{T}v)(t, x)$, where $\hat{u}(s, x) := v_0(s, x)$ for $(s, x) \in [-\tau, 0] \times \mathbb{R}$, $\hat{u}(t, x) := u(t, x)$ for any $(t, x) \in (0, T] \times \mathbb{R}$. Then we have the following two steps.

Step 1: For sufficiently small $T > 0$, $\mathcal{T}(\Gamma_T) \subset \Gamma_T$.

Obviously, $\hat{u} \in C([-\tau, T], \mathcal{Y}^+)$. For any $(t, x) \in (0, T] \times \mathbb{R}$, it follows from (2.1) and $G(v)(t, x) \leq (L + \mu)v(t, x)$ that

$$\begin{aligned} \hat{u}(t, x) &\leq \|v_0(0, \cdot)\|_X e^{-Lt} + e^{-Lt} \frac{1}{L} (e^{Lt} - 1) (L + \mu) K \|v_0\|_{\mathcal{C}_0} \\ &\leq \left(1 + \frac{1}{L} (L + \mu) K (e^{LT} - 1) \right) \|v_0\|_{\mathcal{C}_0} \\ &\leq K \|v_0\|_{\mathcal{C}_0}, \end{aligned}$$

provided that $T > 0$ is sufficiently small such that $(1 + \frac{1}{L} (L + \mu) K (e^{LT} - 1)) \leq K$. Therefore, $\mathcal{T}v \in \Gamma_T$.

Step 2: For sufficiently small $T > 0$, $\mathcal{T} : \Gamma_T \rightarrow \Gamma_T$ is a contraction map.

For any $v_1, v_2 \in \Gamma_T$, define $\hat{u}_1 = \mathcal{T}v_1$ and $\hat{u}_2 = \mathcal{T}v_2$. Then for any $(t, x) \in (0, T] \times \mathbb{R}$, by (2.1), we get

$$\begin{aligned} |\hat{u}_1(t, x) - \hat{u}_2(t, x)| &\leq \int_0^t e^{-L(t-s)} (\Psi(t-s) |G(v_1)(s, \cdot) - G(v_2)(s, \cdot)|)(x) ds \\ &\leq \int_0^t e^{-L(t-s)} \left(L + \mu + (1 + \beta)(K \|v_0\|_{\mathcal{C}_0} + 1) + \frac{\mu}{\beta} \right) \|v_1 - v_2\|_{\mathcal{C}_T} ds \\ &\leq \frac{1}{L} (e^{LT} - 1) \left(L + \mu + (1 + \beta)(K \|v_0\|_{\mathcal{C}_0} + 1) + \frac{\mu}{\beta} \right) \|v_1 - v_2\|_{\mathcal{C}_T}. \end{aligned}$$

Apparently, if T is sufficiently small, then there exists $0 < \rho < 1$ such that

$$\|\hat{u}_1 - \hat{u}_2\|_{\mathcal{C}_T} = \|\mathcal{T}v_1 - \mathcal{T}v_2\|_{\mathcal{C}_T} \leq \rho \|v_1 - v_2\|_{\mathcal{C}_T}, \quad (2.3)$$

which indicates that \mathcal{T} is a contraction map on Γ_T .

By the Banach's fixed point theorem, there exists a unique function $v \in \Gamma_T$, which is the fixed point of \mathcal{T} . Especially, $v \in \Gamma_T$ is the unique mild solution of (1.2) on $t \in (0, T]$ and satisfies

$$v(t, x) := e^{-Lt}(\Psi(t)v_0(\cdot))(x) + \int_0^t e^{-L(t-s)}(\Psi(t-s)G(v)(s, \cdot))(x)ds \quad (2.4)$$

for $(t, x) \in (0, T] \times \mathbb{R}$. This confirms the local existence of mild solution $v(t, x)$ of (1.2). To establish the global existence of mild solution $v(t, x)$ of (1.2), we can assume that $v(t, x)$ is the unique non-continuable mild solution of (1.2) on the maximal existence interval $[0, \sigma_{v_0})$ for

$\sigma_{v_0} > 0$. Apparently, $v \in C([- \tau, \sigma_{v_0}), \mathcal{Y}^+)$. Since $v(t, x)$ satisfies (2.4) for any $t > 0$, it follows that

$$v(t, x) \leq e^{-Lt}(\Psi(t)v_0(\cdot))(x) + \int_0^t e^{-L(t-s)}(\Psi(t-s)(L + \mu)v(s, \cdot))(x)ds$$

for all $(t, x) \in (0, \sigma_{v_0}) \times \mathbb{R}$. Consider the function $v^+(t) := e^{\mu t}\|v_0\|_{C_0}$, where $t \geq 0$. It is obvious that $v^+(t)$ satisfies equation $u_t = \Delta u(t, x) + \mu u(t, x)$ with the initial data $u(0, x) = \|v_0\|_{C_0}$ for any $(t, x) \in (0, +\infty) \times \mathbb{R}$. By the comparison principle (see [16, Proposition 3]), we get that $v(t, x) \leq v^+(t)$ for any $(t, x) \in [0, \sigma_{v_0}) \times \mathbb{R}$. Thus, $\sigma_{v_0} = +\infty$ and

$$v(t, x) \leq e^{\mu t}\|v_0\|_{C_0} \quad (2.5)$$

for any $(t, x) \in [0, +\infty) \times \mathbb{R}$. Furthermore, it is known that the mild solution $v(t, x)$ of (1.2) is a classical solution for $t > \tau$ [16, 24].

This completes the proof. \square

Proof of Theorem 1.2:

Step 1: Proof of (1.5).

Suppose that $v_0(s, x) = 0$ for $|x| \geq l$, $s \in [-\tau, 0]$, where l is a positive constant. Since $v(t, x) \geq 0$ for all $t \geq -\tau$ and $x \in \mathbb{R}$, one has

$$\mu v(t, x) \left(\frac{1 - v(t - \tau, x)}{1 + \beta v(t - \tau, x)} \right) \leq \frac{\mu v(t, x)}{1 + \beta v(t - \tau, x)} \leq \mu v(t, x), \quad (t, x) \in [0, +\infty) \times \mathbb{R}.$$

Suppose $w(t, x)$ with $w(0, x) = v_0(0, x)$ solve the equation $w_t(t, x) = w_{xx}(t, x) + \mu w(t, x)$, $t > 0$, $x \in \mathbb{R}$. It follows from the comparison principle that

$$0 \leq v(t, x) \leq w(t, x) := \frac{e^{\mu t}}{\sqrt{4\pi t}} \int_{-l}^l e^{-(x-y)^2/(4t)} v_0(0, y) dy, \quad \forall t \geq 0, x \in \mathbb{R}. \quad (2.6)$$

Let $c > 2\sqrt{\mu}$, we then obtain

$$0 \leq v(t, x) \leq \frac{e^{\mu t}\|v_0\|_{C_0}}{\sqrt{4\pi t}} \int_{-l}^l e^{-(ct-L)^2/(4t)} dy, \quad \forall t \geq \frac{l}{c}, |x| \geq ct,$$

which further yields (1.5). It is noteworthy that, because $v(t, x)$ is continuous, non-negative and converges to 0 at $\pm\infty$ (see (2.6)), the maximum of $v(t, x)$ on $(-\infty, -ct] \cup [ct, +\infty)$ can be reached.

Step 2: The proof of (1.4).

For any $(t, x) \in [2\tau, \infty) \times \mathbb{R}$, we have

$$\begin{aligned} v_t(t, x) &= v_{xx}(t, x) + \mu v(t, x) \left(\frac{1 - v(t - \tau, x)}{1 + \beta v(t - \tau, x)} \right) \\ &= v_{xx}(t, x) + \mu h(t, x)v(t, x). \end{aligned}$$

Since $h(t, x) := \frac{1-v(t-\tau, x)}{1+\beta v(t-\tau, x)}$ is continuous in $(t, x) \in [2\tau, \infty) \times \mathbb{R}$, $\mu|h(t, x)| \leq \frac{\mu(1+\beta)}{\beta}$ for any $(t, x) \in [2\tau, \infty) \times \mathbb{R}$ and $v(t, x) > 0$ for any $(t, x) \in [2\tau, \infty) \times \mathbb{R}$. In order to apply the Harnack inequalities in [5, Lemma 3.8], we first take $t^* \geq 3\tau$, $x^* \in \mathbb{R}$, then choose $\theta = \frac{\tau}{3}$. Let $\Omega = (x^* - 3, x^* + 3)$, $U = (x^* - 2, x^* + 2)$ and $D = (x^* - 1, x^* + 1)$, in addition, to avoid the confusion, we denote τ in [5, Lemma 3.8] as $\hat{\tau}$, and let $\hat{\tau} = t^* - \frac{4}{3}\tau$, then there exists a constant

$\gamma > 0$ which is independent of t^* and x^* , such that

$$0 < \sup_{[t^*-\tau, t^*-\frac{2\tau}{3}] \times [x^*-1, x^*+1]} v(s, y) \leq \gamma \inf_{[t^*-\frac{\tau}{3}, t^*] \times [x^*-1, x^*+1]} v(s, y).$$

Due to the arbitrariness of t^* and x^* , it holds

$$0 < v(t - \tau, x) \leq \gamma v(t, x), \quad \forall t > 3\tau, \quad x \in \mathbb{R}.$$

Let $g(\varpi) = \frac{1-\varpi}{1+\beta\varpi}$. Since $g(\varpi)$ is decreasing with respect to $\varpi \in [0, \infty)$, we could obtain

$$\begin{aligned} v_t(t, x) &= v_{xx}(t, x) + \mu v(t, x) \left(\frac{1 - v(t - \tau, x)}{1 + \beta v(t - \tau, x)} \right) \\ &\geq v_{xx}(t, x) + \mu v(t, x) \left(\frac{1 - \gamma v(t, x)}{1 + \beta \gamma v(t, x)} \right), \quad \forall t > 3\tau, \quad x \in \mathbb{R}. \end{aligned}$$

Let $z(t, x)$ with $z(3\tau, x) = v(3\tau, x)$ solve

$$z_t(t, x) = z_{xx}(t, x) + \mu z(t, x) \left(\frac{1 - \gamma z(t, x)}{1 + \beta \gamma z(t, x)} \right), \quad \forall t > 3\tau, \quad x \in \mathbb{R}. \quad (2.7)$$

Clearly, $v(t, x)$ is a upper solution for equation (2.7). Besides, equation (2.7) admits two equilibria $z = 0$ and $z = \gamma^{-1}$. It follows from the classical theory of asymptotic spreading in [1, §4] that the solution of (2.7) satisfies

$$\liminf_{t \rightarrow \infty} \min_{|x| \leq ct} z(t, x) \rightarrow \gamma^{-1} \quad \text{for all } 0 \leq c < 2\sqrt{\mu}.$$

Then by the comparison principle, we get

$$\liminf_{t \rightarrow \infty} \min_{|x| \leq ct} v(t, x) \geq \limsup_{t \rightarrow \infty} \max_{|x| \leq ct} z(t, x) \geq \liminf_{t \rightarrow \infty} \min_{|x| \leq ct} z(t, x) = \gamma^{-1}$$

for any $0 \leq c < 2\sqrt{\mu}$.

This completes the proof. \square

§3 Numerical simulations

In this section, the numerical simulations are demonstrated by the finite difference scheme. Precisely, we consider the case of a finite spatial domain with homogeneous Neumann boundary condition, which models a closed environment with reflecting boundaries, i.e., the individual cannot leave the domain. For equation (1.2), we choose $\mu = 1$, $x \in [-100, 100] := \Omega$, $t \in [0, 250]$, that is,

$$\begin{cases} v_t(t, x) = v_{xx}(t, x) + v(t, x) \left(\frac{1 - v(t - \tau, x)}{1 + \beta v(t - \tau, x)} \right), \\ v_x(-100, t) = v_x(100, t) = 0, \\ v(s, x) = v_0(s, x), \quad s \in [-\tau, 0]. \end{cases} \quad (3.1)$$

The initial condition is

$$v_0(s, x) = \begin{cases} 1, & x \in [-\alpha, \alpha], \\ 0, & x \in \Omega \setminus [-\alpha, \alpha], \end{cases}$$

where $\alpha > 0$. We then use the finite difference method to discretize (3.1). Define the uniform partition of the domain $\Omega = [-100, 100]$ by:

$$-100 = x_1 < x_2 < x_3 < \cdots < x_{n-1} < x_n < x_{n+1} = 100,$$

where $h = 200/n = 0.2$ and $x_i = x_1 + (i - 1) \cdot h$, $i = 1, 2, \dots, n + 1$. The time domain $[0, 250]$ is treated as $k = 250/m = 0.02$, $t_j = t_1 + (j - 1) \cdot k$, $j = 1, 2, \dots, m + 1$. The following are numerical simulations:




Figure 1: Choose $\alpha = 20$, $\beta = 0.5$ and $\tau = 0.2$. The left is the 3D-graph of $v(t, x)$ and the right is the graph of $v(t, 100)$. The solution approaches positive steady states.




Figure 2: Choose $\alpha = 20$, $\beta = 0.5$ and $\tau = 1.8$. The left is the 3D-graph of $v(t, x)$ and the right is the graph of $v(t, 100)$. The solution approaches positive steady states but with oscillations.




Figure 3: Choose $\alpha = 20$, $\beta = 0.5$ and $\tau = 2.4$. The left is the 3D-graph of $v(t, x)$ and the right is the graph of $v(t, 100)$. The solution is oscillatory and eventually converges to a time-periodic solution.

According to the figures given above, as time increases, for small τ , the nonnegative steady states for (3.1) are $v \equiv 0$ and $v \equiv 1$; for sufficiently large τ , the steady $v \equiv 1$ becomes unstable, the non-constant bounded steady states may exist. In particular, if we choose $\tau = 0.2$ and $\tau = 1.8$, the solution finally connects the equilibria 0 to 1. When choosing $\tau = 2.4$, as time increases, there exists an oscillating solution, but the solution still has upper and lower bounds and it seems to be a spatially uniform temporally periodic solution. Note that, for the rigorous

proof of the existence of the time periodic solution of the delayed food-limited model, readers can refer to [20,21,27]. Especially, Su et al. [20,21] investigated the spatial structure of system (3.1) with Dirichlet boundary conditions, and showed that the solution tends to a stable spatial nonhomogeneous time-periodic solution when delay τ is large enough.

Moreover, the simulations support our theoretical results, i.e., Theorem 1.2. As we can see from figure 1 and 2, there hold

$$(i) \quad \lim_{t \rightarrow +\infty, |x| \geq ct} v(t, x) = 0, \forall c > 2;$$

$$(ii) \quad \liminf_{t \rightarrow +\infty, |x| \leq ct} v(t, x) = 1, \forall 0 < c < 2.$$

In figure 3, there exists a constant $\kappa > 0$ such that

$$(i) \quad \lim_{t \rightarrow +\infty, |x| \geq ct} v(t, x) = 0, \forall c > 2;$$

$$(ii) \quad \liminf_{t \rightarrow +\infty, |x| \leq ct} v(t, x) > \kappa, \forall 0 < c < 2.$$

It follows from these simulations that the spreading speed c^* of solution of (3.1) is equal to 2, which is the same as the conclusion of Theorem 1.2.

Acknowledgement

The authors are grateful to the referees for their valuable suggestions that help improvement of the manuscript. The authors would also like to thank Prof. Zhicheng Wang for his insightful comments and suggestions.

Declarations

Conflict of interest The authors declare no conflict of interest.

References

- [1] D G Aronson, H F Weinberger. *Nonlinear diffusion in population genetics, combustion, and nerve pulse propagation*, in: J A Goldstein (Ed.), *Partial Differential Equations and Related Topics*, in: *Lecture Notes in Mathematics*, Springer, Berlin, 1975, 446: 5-49.
- [2] D Daners, P K Medina. *Abstract Evolution Equations, Periodic Problems and Applications*, Pitman Research Notes in Mathematics Series, Longman, Harlow, UK, 1992, 279.
- [3] W Feng, X Lu. *On diffusive population models with toxicants and time delays*, *J Math Anal Appl*, 1999, 233: 373-386.
- [4] W Feng, X Lu. *Global periodicity in a class of reaction-diffusion systems with time delays*, *Discrete Cont Dyn Syst*, 2003, 3B: 69-78.

- [5] J Földes, P Poláčik. *On cooperative parabolic systems: Harnack inequalities and asymptotic symmetry*, Discrete Contin Dyn Syst, 2009, 25(1): 133-157.
- [6] K Gopalsamy, M R S Kulenovic, G Ladas. *Time lags in a food-limited population model*, Appl Anal, 1988, 31: 225-237.
- [7] K Gopalsamy, M R S Kulenovic, G Ladas. *Environmental periodicity and time delays in a food-limited population model*, J Math Anal Appl, 1990, 147: 545-555.
- [8] S A Gourley, M A J Chaplain. *Travelling fronts in a food-limited population model with time delay*, Proc R Soc Edinburgh Sect A, 2002, 132: 75-89.
- [9] S A Gourley, J W H So. *Dynamics of a food-limited population model incorporating nonlocal delays on a finite domain*, J Math Biol, 2002, 44: 49-78.
- [10] G E Hutchinson. *Circular causal systems in ecology*, Ann N Y Acad Sci, 1948, 50: 221-246.
- [11] F Hamel, L Ryzhik. *On the nonlocal Fisher-KPP equation: steady states, spreading speed and global bounds*, Nonlinearity, 2014, 27: 2735-3753.
- [12] K Hasik, J Kopfová, P Nábělková, S Trofimchuk. *On the Geometric Diversity of Wavefronts for the Scalar Kolmogorov Ecological Equation*, J Nonlinear Sci, 2020, <https://doi.org/10.1007/s00332-020-09642-9>
- [13] Y Kuang. *Delay Differential Equations with Applications in Population Dynamics*, Academic Press, New York, 1993.
- [14] G Lin. *Spreading speed of the delayed Fisher equation without quasimonotonicity*, Nonlinear Anal Real World Appl, 2011, 12(6): 3713-3718.
- [15] X L Liu, S X Pan. *Spreading speed in a nonmonotone equation with dispersal and delay*, Mathematics, 2019, 7(3): 291, <https://doi.org/10.3390/math7030291>.
- [16] R H Martin, H L Smith. *Abstract functional differential equations and reaction-diffusion systems*, Trans Am Math Soc, 1990, 321: 1-44.
- [17] E C Pielou. *An Introduction to Mathematical Ecology*, Wiley, New York, 1969.
- [18] F E Smith. *Population dynamics in Daphnia magna*, Ecology, 1963, 44: 651-663.
- [19] J W H So, J S Yu. *On the uniform stability for a food-limited population model with time delay*, Proc R Soc Edinburgh Sect A, 1995, 125: 991-1005.
- [20] Y Su, A Y Wan, J J Wei. *Bifurcation analysis in a diffusive 'food-limited' model with time delay*, Appl Anal, 2010, 89(7): 1161-1181.
- [21] Y Su, J J Wei, J P Shi. *Hopf bifurcations in a reaction-diffusion population model with delay effect*, J Differential Equations, 2009, 247(4): 1156-1184.
- [22] E Trofimchuk, M Pinto, S Trofimchuk. *Existence and uniqueness of monotone wavefronts in a nonlocal resource-limited model*, Proceedings of the Royal Society of Edinburgh: Section A Mathematics, 2020, 150: 2462-2483.
- [23] G Tian, Z C Wang. *Spreading speed in a food-limited population model with nonlocal delay*, Appl Math Lett, 2020, 102: 106121.

- [24] J Wu. *Theory and Applications of Partial Functional Differential Equations*, Springer-Verlag, New York, 1996.
- [25] Z C Wang, W T Li. *Monotone travelling fronts of a food-limited population model with nonlocal delay*, Nonlinear Analysis: Real World Applications, 2007, 18: 699-712.
- [26] Z C Wang, W T Li, S Ruan. *Travelling wave fronts of reaction-diffusion systems with spatio-temporal delays*, J Differential Equations, 2006, 222: 185-232.
- [27] A Y Wan, J J Wei. *Hopf bifurcation analysis of a food-limited population model with delay*, Nonlinear Anal Real World Appl, 2010, 11(2): 1087-1095.
- [28] J D Wei, L X Tian, J B Zhou, Z L Zhen, J Xu. *Existence and asymptotic behavior of traveling wave fronts for a food-limited population model with spatio-temporal delay*, Japan J Indust Appl Math, 2017, 34: 305-320.

¹College of Mathematics and Statistics, Northwest Normal University, Lanzhou 730070, China.

Email: tiang@nwnu.edu.cn

²School of Mathematics and Statistics, Lanzhou University, Lanzhou 730000, China.

Email: tiang17@lzu.edu.cn

³School of Mathematics, China University of Mining and Technology, Xuzhou 221008, China.

Email: an18020536372@163.com