

On some properties of the bibasic Humbert hypergeometric functions Ξ_1 and Ξ_2

CAI Qing-bo^{1,*} Ghazi S. Khammash²
Shimaa I. Moustafa³ Ayman Shehata^{3,4}

Abstract. The main object of this paper is to deduce the bibasic Humbert functions Ξ_1 and Ξ_2 . Some interesting results and elementary summations technique that was successfully employed, q -recursion, q -derivatives relations, the q -differential recursion relations, the q -integral representations for Ξ_1 and Ξ_2 are given. The summation formula derives a list of p -analogues of transformation formulas for bibasic Humbert functions that have been studied, also some hypergeometric functions properties of some new interesting special cases have been given.

§1 Introduction

The bibasic summation formula and multivariable hypergeometric functions have been investigated and have a long history see, for example, [3, 11, 13, 36], also for two variables or more multivariable basic hypergeometric functions see, for example, [1, 16, 21, 29, 33, 40]. Recently, considerable attention from several researchers and mathematicians has been developed and studied the bibasic hypergeometric functions in [2, 4, 14, 20, 23, 29] were presented, the basic hypergeometric functions, which namely q -analogue generalizations of the ordinary hypergeometric series have a long history [23]. The derivatives, q -derivatives of the generalization of the hypergeometric series and basic hypergeometric series with respect to the parameters see, also for example, [14, 24–26]. Bibasic q -Appell functions with contiguous bibasic q -Appell functions, q -statistical summability method [17] and certain q -partial derivative relations have been derived in [36]. Also, the authors [36, 37] introduce some summation formulas for the bibasic q -Appell function and the relation between the bibasic Appell series and continued fractions of Ramanujan. Some multivariable generalizations of the bibasic summation formula have been studied in [3, 11, 22]. Also, some transformation formulas expressing certain sums of bibasic series by using contour integrals and the calculus of residues defined by Agarwal and Verma, see [11], for q -calculus and q -analogue, see some examples as [5, 6, 8, 9, 18]. Some extensions of the bibasic indefinite summation formula to derive bibasic extensions of Euler's transformation formula and of fields and Wimp expansion formula have been obtained by Gasper [13].

Received: 2022-09-14. Revised: 2023-05-07.

MR Subject Classification: 33D65, 05A30, 33D70, 33C65, 33C70.

Keywords: q -calculus, bibasic Humbert hypergeometric functions, q -derivative.

Digital Object Identifier(DOI): <https://doi.org/10.1007/s11766-023-4848-8>.

Supported by the National Natural Science Foundation of China(11601266) and the Natural Science Foundation of Fujian Province of China(2020J01783).

*Corresponding author.

Because of their utility and applications in a variety of research fields, the bibasic hypergeometric functions associated with special matrix functions, matrix series have received recent attention see, e.g., [4, 31, 32], and also see, e.g., [10, 19, 27, 28, 30, 34, 35, 39] for some q -function of different type of matrix functions.

The q -analogue of τ , also known as the q -number or q -bracket number $[\tau]_q$ is defined by [12]

$$[\tau]_q = 1 + q + q^2 + \dots + q^{\tau-1} = \frac{1 - q^\tau}{1 - q}; \tau \in \mathbb{C}. \quad (1)$$

For r the factorial $[r]_q!$ is defined by [12]

$$[r]_q! = \begin{cases} [1]_q + [2]_q + \dots + [r]_q, & r \in \mathbb{N}; \\ 1, & r = 0. \end{cases} \quad (2)$$

The q -shifted factorial is defined by [12]

$$(q^x; q)_\tau = \begin{cases} \prod_{r=0}^{\tau-1} (1 - q^{x+r}), & \tau \geq 1; \\ 1, & \tau = 0. \end{cases} \quad (3)$$

$$= \begin{cases} (1 - q^x)(1 - q^{x+1}) \dots (1 - q^{x+\tau-1}), & \tau \in \mathbb{N}, q^x \in \mathbb{C} \setminus \{1, 1/q, 1/q^2, \dots, 1/q^{\tau-1}\}; \\ 1, & x \in \mathbb{C}, 0 < |q| < 1, q \in \mathbb{C} - \{1\}. \end{cases}$$

The p -analogue of gamma functions $\Gamma_p(c)$ is defined by the q -integral representations as follows

$$\Gamma_p(c) = \int_0^{\frac{1}{1-p}} E_p(-pu) u^{c-1} d_p u, \quad (4)$$

where

$$E_p(x) = \sum_{\tau=0}^{\infty} \frac{p^{\binom{\tau}{2}}}{[\tau]_p!} x^\tau.$$

The q -binomial theorem is

$$\phi_1(p^a; -, p, x) = \frac{(xp^a; p)_\infty}{(x; p)_\infty}, |x| < 1. \quad (5)$$

In [12], the basic or q -hypergeometric function has been defined as follows

$$\phi_1(p^a, p^b; p^c; p, x) = \sum_{\tau=0}^{\infty} \frac{(p^a; p)_\tau (p^b; p)_\tau}{(p^c; p)_\tau (p; p)_\tau} x^\tau, p^c \neq 1, q^{-1}, q^{-2}, \dots, |x| < 1. \quad (6)$$

Now we may define a bibasic Humbert functions Ξ_1 and Ξ_2 on the two independent bases p and q , for $0 < |p| < 1$, $0 < |q| < 1$, $p, q \in \mathbb{C}$, as follows:

$$\begin{aligned} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\ = \sum_{k, \ell=0}^{\infty} \frac{(p^a; p)_k (q^b; q)_\ell (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^\ell y^k, p^d \neq 1, q^{-1}, q^{-2}, \dots, |x| < 1, |y| < 1 \end{aligned} \quad (7)$$

and

$$\begin{aligned} \Xi_2(p^a, q^b; p^c; p, q, x, y) \\ = \sum_{k, \ell=0}^{\infty} \frac{(p^a; p)_k (q^b; q)_\ell}{(p^c; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^\ell y^k, p^c \neq 1, q^{-1}, q^{-2}, \dots, |x| < 1, |y| < 1. \end{aligned} \quad (8)$$

Also, q -difference operator defined by [15]

$$D_{y,q} f(y) = [f(y) - f(yq)] [(1 - q)y]^{-1}, \quad (9)$$

where f is a function real or complex -valued and $D_{y,q} f(0) = f'(0)$ provided $f(0)$ exists.

§2 Some new auxiliary results

Here we utilize the interesting results and elementary summations technique that was successfully employed in the bibasic Humbert confluent hypergeometric functions Ξ_1 and Ξ_2 on two independent bases p and q of two variables.

Theorem 2.1. *The following q -recursion relations for Ξ_1 and Ξ_2 are true*

$$\begin{aligned} \Xi_1(p^{a+1}, q^b, p^c; p^d; p, q, x, y) = & \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\ & + \frac{p^a(1-p^c)y}{1-p^d} \Xi_1(p^{a+1}, q^b, p^{c+1}; p^{d+1}; q, p, x, y), p^d \neq 1, \end{aligned} \quad (10)$$

$$\begin{aligned} \Xi_1(p^a, q^b, p^{c+1}; p^d; q, p, x, y) = & \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\ & + \frac{p^c(1-p^a)y}{1-p^d} \Xi_1(p^{a+1}, q^b, p^{c+1}; p^{d+1}; q, p, x, y), q^d \neq 1, \\ \Xi_1(p^a, p^b, p^c; p^{d-1}; q, p, x, y) = & \frac{1}{1-p^{d-1}} \Xi_1(p^a, q^b, p^c; p^d; q, p, x, y) - \frac{p^{d-1}}{1-p^{d-1}} \Xi_1(p^a, q^b, p^c; p^d; q, p, px, y) \\ & + \frac{p^{d-1}(1-p^a)(1-p^c)y}{(1-p^{d-1})(1-p^d)} \Xi_1(p^{a+1}, q^b, p^{c+1}; p^{d+1}; q, p, px, y), p^d, p^{d-1} \neq 1, \end{aligned} \quad (11)$$

$$\begin{aligned} = & \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) + \frac{p^{d-1}(1-p^a)(1-p^c)y}{(1-p^{d-1})(1-p^d)} \Xi_1(p^{a+1}, q^b, p^{c+1}; p^{d+1}; q, p, x, y) \\ & + \frac{p^{d-1}}{1-p^{d-1}} \Xi_1(p^a, q^b, p^c; p^d; q, p, x, py) - \frac{p^{d-1}}{1-p^{d-1}} \Xi_1(p^a, q^b, p^c; p^d; q, p, px, py), p^d, p^{d-1} \neq 1 \end{aligned}$$

and

$$\begin{aligned} \Xi_2(p^{a+1}, q^b; p^c; p, q, x, y) = & \Xi_2(p^a, q^b; p^c; p, q, x, y) + \frac{p^a y}{1-p^c} \Xi_2(p^{a+1}, q^b; p^{c+1}; q, p, x, y), p^c \neq 1, \\ \Xi_2(p^a, p^b; p^{c-1}; q, p, x, y) = & \frac{1}{1-p^{c-1}} \Xi_2(p^a, q^b; p^c; q, p, x, y) - \frac{p^{c-1}}{1-p^{c-1}} \Xi_2(p^a, q^b; p^c; q, p, px, y) \\ & + \frac{p^{c-1}(1-p^a)y}{(1-p^{c-1})(1-p^c)} \Xi_2(p^{a+1}, q^b; p^{c+1}; q, p, px, y), p^c, p^{c-1} \neq 1, \\ = & \Xi_2(p^a, q^b; p^c; p, q, x, y) + \frac{p^{c-1}(1-p^a)y}{(1-p^{c-1})(1-p^c)} \Xi_2(p^{a+1}, q^b; p^{c+1}; q, p, x, y) \\ & + \frac{p^{c-1}}{1-p^{c-1}} \Xi_2(p^a, q^b; p^c; q, p, x, py) - \frac{p^{c-1}}{1-p^{c-1}} \Xi_2(p^a, q^b; p^c; q, p, px, py), p^c, p^{c-1} \neq 1. \end{aligned} \quad (12)$$

Proof. To prove (10), we proceed as follows

$$(p^{a+1}; p)_k = \frac{1-p^{a+k}}{1-p^a} (p^a; p)_k = \left[1 + p^a \frac{1-p^k}{1-p^a} \right] (p^a; p)_k,$$

and

$$\begin{aligned} (p^a; p)_{k+1} &= (1-p^a)(p^{a+1}; p)_k, \\ (p^c; p)_{k+1} &= (1-p^c)(p^{c+1}; p)_k, \\ (p^d; p)_{\ell+k+1} &= (1-p^d)(p^{d+1}; p)_{\ell+k}, \end{aligned}$$

we have

$$\begin{aligned}
 & \Xi_1(p^{a+1}, q^b, p^c; p^d; p, q, x, y) - \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\
 &= p^a \sum_{\ell, k=0}^{\infty} \left[\frac{1-p^k}{1-p^a} \right] \frac{(p^a; p)_k (q^b; q)_\ell (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^\ell y^k \\
 &= \frac{p^a}{1-p^a} \sum_{\ell=0, k=1}^{\infty} \frac{(p^a; p)_k (q^b; q)_\ell (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_{k-1} (q; q)_\ell} x^\ell y^k \\
 &= \frac{p^a}{1-p^a} \sum_{\ell, k=0}^{\infty} \frac{(p^a; p)_{k+1} (q^b; q)_\ell (p^c; p)_{k+1}}{(p^d; p)_{\ell+k+1} (p; p)_k (q; q)_\ell} x^\ell y^{k+1} \\
 &= \frac{p^a(1-p^c)y}{1-p^d} \Xi_1(p^{a+1}, p^b, p^{c+1}; p^{d+1}; q, p, x, y).
 \end{aligned}$$

Similarly, by same manner the relations (11)-(12) can be obtained. $\square$

Theorem 2.2. *The following relations for Ξ_1 and Ξ_2 are true*

$$\begin{aligned}
 & \Xi_1(p^a, q^{b+1}, p^c; p^d; q, p, x, y) = \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\
 & + \frac{q^b x}{1-p^d} \Xi_1(p^a, q^{b+1}, p^c; p^{d+1}; q, p, x, y), p^d \neq 1,
 \end{aligned} \tag{13}$$

$$\begin{aligned}
 & (1-q^b) \Xi_1(p^a, q^{b+1}, p^c; p^d; q, p, x, y) + q^b \Xi_1(p^a, q^b, p^c; p^d; q, p, qx, y) \\
 &= \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y), \\
 & \Xi_1(p^a, q^{b+1}, p^c; p^d; q, p, x, y) \\
 &= \Xi_1(p^a, q^b, p^c; p^d; q, p, qx, y) + \frac{x}{1-p^d} \Xi_1(p^a, q^{b+1}, p^c; p^{d+1}; q, p, x, y), p^d \neq 1
 \end{aligned} \tag{14}$$

and

$$\begin{aligned}
 & (1-q^b) \Xi_2(p^a, q^{b+1}, p^c; q, p, x, y) + q^b \Xi_2(p^a, q^b, p^c; q, p, qx, y) = \Xi_2(p^a, q^b, p^c; p, q, x, y), \\
 & \Xi_2(p^a, q^{b+1}, p^c; q, p, x, y) \\
 &= \Xi_2(p^a, q^b, p^c; p, q, x, y) + \frac{q^b x}{1-p^c} \Xi_2(p^a, q^{b+1}, p^{c+1}; q, p, x, y), p^c \neq 1, \\
 & \Xi_2(p^a, q^{b+1}, p^c; q, p, x, y) \\
 &= \Xi_2(p^a, q^b, p^c; q, p, qx, y) + \frac{x}{1-p^c} \Xi_2(p^a, q^{b+1}, p^{c+1}; q, p, x, y), p^c \neq 1.
 \end{aligned} \tag{15}$$

Proof. Using the relation

$$(q^b; q)_{\ell+1} = (1-q^b)(q^{b+1}; q)_\ell = (1-q^{b+\ell})(q^b; q)_\ell,$$

we have

$$\begin{aligned}
 & \Xi_1(p^a, q^{b+1}, p^c; p^d; q, p, x, y) - \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\
 &= q^b \sum_{\ell, k=0}^{\infty} \left[\frac{1-q^\ell}{1-q^b} \right] \frac{(p^a; p)_k (q^b; q)_\ell (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^\ell y^k \\
 &= \frac{q^b}{1-q^b} \sum_{\ell=1, k=0}^{\infty} \frac{(p^a; p)_k (q^b; q)_\ell (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_k (q; q)_{\ell-1}} x^\ell y^k \\
 &= \frac{q^b x}{1-q^b} \sum_{\ell, k=0}^{\infty} \frac{(p^a; p)_k (q^b; q)_{\ell+1} (p^c; p)_k}{(p^d; p)_{\ell+k+1} (p; p)_k (q; q)_\ell} x^\ell y^k.
 \end{aligned}$$

Now, by making use of the relation (7), we obtain

$$= \frac{q^b x}{1 - p^d} \Xi_1(p^a, q^{b+1}, p^c; p^{d+1}; q, p, x, y), p^d \neq 1.$$

Similarly, we can easily prove the subsequent results (14)-(15). $\square$

Theorem 2.3. The q -derivatives relations for Ξ_1 and Ξ_2 holds true

$$D_{x,q}^m \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) = \frac{(q^b; q)_m}{(1 - q)^m (p^d; p)_m} \Xi_1(p^a, q^{b+m}, p^c; p^{d+m}; q, p, x, y), \quad (16)$$

$$D_{y,p}^n \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) = \frac{(p^a; p)_n (p^c; p)_n}{(1 - p)^n (p^d; p)_n} \Xi_1(p^{a+n}, q^b, p^{c+n}; p^{d+n}; q, p, x, y),$$

$$\begin{aligned} D_{x,p}^m D_{y,p}^n \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\ = \frac{(p^a; p)_n (q^b; q)_m (p^c; p)_n}{(1 - q)^m (1 - p)^n (p^d; p)_{m+n}} \Xi_1(p^{a+n}, q^{b+m}, p^{c+n}; p^{d+m+n}; q, p, x, y) \end{aligned} \quad (17)$$

and

$$\begin{aligned} D_{x,q}^m \Xi_2(p^a, q^b; p^c; p, q, x, y) &= \frac{(q^b; q)_m}{(1 - q)^m (q^c; q)_m} \Xi_2(q^a, q^{b+m}; p^{c+m}; q, p, x, y), \\ D_{y,p}^n \Xi_2(p^a, q^b; p^c; p, q, x, y) &= \frac{(p^a; p)_n}{(1 - p)^n (p^c; p)_n} \Xi_2(q^{a+n}, p^b; p^{c+n}; q, p, x, y), \end{aligned} \quad (18)$$

$$\begin{aligned} D_{x,q}^m D_{y,p}^n \Xi_2(p^a, q^b; p^c; p, q, x, y) \\ = \frac{(p^a; p)_n (q^b; q)_m}{(1 - q)^m (1 - p)^n (p^c; p)_{m+n}} \Xi_2(q^{a+n}, p^{b+m}; q^{c+m+n}; q, p, x, y). \end{aligned}$$

Proof. From q -derivative in (9), we yield

$$\begin{aligned} D_{x,q} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\ = \sum_{\ell, k=0}^{\infty} \frac{1 - q^\ell}{1 - q} \frac{(p^a; p)_k (q^b; q)_\ell (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^{\ell-1} y^k \\ = \sum_{\ell, k=0}^{\infty} \frac{(p^a; p)_k (q^b; q)_{\ell+1} (p^c; p)_k (1 - q)^{-1}}{(p^d; p)_{\ell+k+1} (p; p)_k (q; q)_\ell} x^\ell y^k \\ = \frac{(1 - q^b)}{(1 - q)(1 - p^d)} \sum_{\ell, k=0}^{\infty} \frac{(p^a; p)_k (q^{b+1}; q)_\ell (p^c; p)_k}{(p^{d+1}; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^\ell y^k \\ = \frac{(1 - q^b)}{(1 - q)(1 - p^d)} \Xi_1(p^a, q^{b+1}, p^c; p^{d+1}; q, p, x, y) \end{aligned} \quad (19)$$

and

$$\begin{aligned} D_{y,p} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\ = \sum_{\ell=0, k=1}^{\infty} \frac{1}{1 - p} \frac{(p^a; p)_k (q^b; q)_\ell (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_{k-1} (q; q)_\ell} x^\ell y^{k-1} \\ = \frac{(1 - p^a)(1 - p^c)}{(1 - p)(1 - p^d)} \sum_{\ell, k=0}^{\infty} \frac{(p^{a+1}; p)_k (q^b; q)_\ell (p^{c+1}; p)_k}{(p^{d+1}; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^\ell y^k \\ = \frac{(1 - p^a)(1 - p^c)}{(1 - p)(1 - p^d)} \Xi_1(p^{a+1}, q^b, p^{c+1}; p^{d+1}; q, p, x, y). \end{aligned} \quad (20)$$

Now by iterating q -derivative on Ξ_1 for m -times and n -times, we get (16) and (17). By applying the relation of the q -derivative in (8), we obtained (18). $\square$

Theorem 2.4. *The recursion relations of q -differential for Ξ_1 and Ξ_2 holds true:*

$$xD_{x,q}\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) = \frac{1-q^b}{(1-q)q^b} \left[\Xi_1(p^a, q^{b+1}, p^c; p^d; q, p, x, y) - \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \right], \quad (21)$$

$$\begin{aligned} xD_{x,q}\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) &= \frac{1-q^b}{1-q} \left[\Xi_1(p^a, q^{b+1}, p^c; p^d; q, p, x, y) - \Xi_1(p^a, q^b, p^c; p^d; p, q, qx, y) \right], \\ yD_{y,p}\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) &= \frac{1-p^a}{(1-p)p^a} \left[\Xi_1(p^{a+1}, q^b, p^c; p^d; q, p, x, y) - \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \right], \\ yD_{y,p}\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) &= \frac{1-p^c}{(1-p)p^c} \left[\Xi_1(p^a, q^b, p^{c+1}; p^d; q, p, x, y) - \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \right] \end{aligned} \quad (22)$$

and

$$\begin{aligned} xD_{x,q}\Xi_2(p^a, q^b; p^c; p, q, x, y) &= \frac{1-q^b}{(1-q)q^b} \left[\Xi_2(p^a, q^{b+1}; p^c; q, p, x, y) - \Xi_2(p^a, q^b; p^c; p, q, x, y) \right], \\ xD_{x,q}\Xi_2(p^a, q^b; p^c; p, q, x, y) &= \frac{1-q^p}{1-q} \left[\Xi_2(p^a, q^{b+1}; p^c; q, p, x, y) - \Xi_2(p^a, q^b; p^c; p, q, qx, y) \right], \\ yD_{y,p}\Xi_2(p^a, q^b; p^c; p, q, x, y) &= \frac{1-p^a}{(1-p)p^a} \left[\Xi_2(p^{a+1}, q^b; p^c; q, p, x, y) - \Xi_2(p^a, q^b; p^c; p, q, x, y) \right]. \end{aligned} \quad (23)$$

Proof. In view of equation (12) and (13), we get (21). In the same way we can also be obtained (22)-(23). $\square$

Corollary 2.5. *The following relations for Ξ_1 and Ξ_2 holds true:*

$$\begin{aligned} (1-p^{d-1})\Xi_1(p^a, q^b, p^c; q^{d-1}; q, p, x, y) &= (1-p^{d-1})\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\ &+ (1-p)p^{d-1}yD_{y,p}\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) + p^{d-1}\Xi_1(p^a, q^b, p^c; q^d; q, p, x, py) \\ &- p^{d-1}\Xi_1(p^a, q^b, p^c; q^d; q, p, px, py), \\ (1-p^{d-1})\Xi_1(p^a, q^b, p^c; q^{d-1}; q, p, x, y) \end{aligned} \quad (24)$$

$$\begin{aligned}
&= \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) - p^{d-1} \Xi_1(p^a, q^b, p^c; p^d; p, q, px, y) \\
&\quad + (1-p)p^{d-1} y D_{y,p} \Xi_1(p^a, q^b, p^c; p^d; p, q, px, y), \\
(1-p^a) \Xi_1(p^{a+1}, q^b, p^c; p^d; q, p, x, y) &= (1-p) y D_{y,p} \Xi_1(p^a, q^b, p^c; p^d; q, p, x, y) \\
&\quad + (1-p^a) \Xi_1(p^a, q^b, p^c; p^d; q, p, x, py), \\
(1-p^c) \Xi_1(p^a, q^b, p^{c+1}; p^d; q, p, x, y) &= (1-p) y D_{y,p} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\
&\quad + (1-p^c) \Xi_1(p^a, q^b, p^c; p^d; q, p, x, py), \\
(1-q^b) \Xi_1(p^a, q^{b+1}, p^c; p^d; q, p, x, y) &= (1-q) x D_{x,p} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\
&\quad + (1-q^b) \Xi_1(p^a, q^b, p^c; p^d; q, p, qx, y)
\end{aligned} \tag{25}$$

and

$$\begin{aligned}
(1-p^{c-1}) \Xi_2(p^a, q^b; p^{c-1}; q, p, x, y) &= (1-p^{c-1}) \Xi_2(p^a, q^b; p^c; p, q, x, y) + (1-p)p^{c-1} y D_{y,p} \Xi_2(p^a, q^b; p^c; p, q, x, y) \\
&\quad + p^{c-1} \Xi_2(p^a, q^b; p^c; q, p, x, py) - p^{c-1} \Xi_2(p^a, q^b; p^c, q^d; q, p, px, py), \\
(1-p^{c-1}) \Xi_2(p^a, q^b; q^{c-1}; q, p, x, y) &= \Xi_2(p^a, q^b; p^c; p, q, x, y) \\
&\quad - p^{c-1} \Xi_2(p^a, q^b; p^c; p, q, px, y) + (1-p)p^{c-1} y D_{y,p} \Xi_2(p^a, q^b; p^c; p, q, px, y), \\
(1-p^a) \Xi_2(p^{a+1}, q^b; p^c; q, p, x, y) &= (1-p) y D_{y,p} \Xi_2(p^a, q^b; p^c; q, p, x, y) + (1-p^a) \Xi_2(p^a, q^b; p^c, p^c; q, p, x, py), \\
(1-q^b) \Xi_2(p^a, q^{b+1}; p^c; q, p, x, y) &= (1-q) x D_{x,p} \Xi_2(p^a, q^b; p^c; p, q, x, y) + (1-q^b) \Xi_2(p^a, q^b; p^c; q, p, qx, y).
\end{aligned} \tag{26}$$

Proof. In view of (11) and (20) yields (24). Similarly, by same manner the relations (25)-(26) can be obtained. $\square$

Theorem 2.6. The following relations for Ξ_1 and Ξ_2 holds true:

$$\begin{aligned}
D_{d,q} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) &= \frac{1}{1-p^d} \left[y D_{y,q} \Xi_1(p^{d+1}) + \frac{1}{1-p} \Xi_1(p^{d+1}, py) \right. \\
&\quad \left. - \frac{1}{1-p} \Xi_1(p^{d+1}, px, py) \right],
\end{aligned} \tag{27}$$

$$\begin{aligned}
D_{a,q} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) &= -\frac{1}{1-p^a} y D_{y,p} \Xi_1, \\
D_{b,p} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) &= -\frac{1}{1-q^b} x D_{x,q} \Xi_1, \\
D_{c,p} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) &= -\frac{1}{1-p^c} y D_{y,p} \Xi_1, \\
D_{d,q} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) &= \frac{1}{1-p^d} \left[y D_{y,q} \Xi_1(p^{d+1}, px) + \frac{1}{1-p} \Xi_1(p^{d+1}) \right. \\
&\quad \left. - \frac{1}{1-p} \Xi_1(p^{d+1}, px) \right]
\end{aligned} \tag{28}$$

and

$$D_{a,q} \Xi_2(p^a, q^b; p^c; p, q, x, y) = -\frac{1}{1-p^a} y D_{y,p} \Xi_2,$$

$$\begin{aligned}
D_{b,p}\Xi_2(p^a, q^b; p^c; p, q, x, y) &= -\frac{1}{1-q^b}x D_{x,q}\Xi_2, \\
D_{c,q}\Xi_2(p^a, q^b; p^c; p, q, x, y) &= \frac{1}{1-p^c} \left[y D_{y,q}\Xi_2(p^{c+1}) + \frac{1}{1-p}\Xi_2(p^{c+1}, py) \right. \\
&\quad \left. - \frac{1}{1-p}\Xi_2(p^{c+1}, px, py) \right], \\
D_{c,q}\Xi_2(p^a, q^b; p^c; p, q, x, y) &= \frac{1}{1-p^c} \left[y D_{y,q}\Xi_2(p^{c+1}, px) + \frac{1}{1-p}\Xi_2(p^{c+1}) \right. \\
&\quad \left. - \frac{1}{1-p}\Xi_2(p^{c+1}, px) \right].
\end{aligned} \tag{29}$$

Proof. Now we have

$$\begin{aligned}
D_{d,p}\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) &= \sum_{\ell, k=0}^{\infty} \left[\frac{1}{(p^d; p)_k} - \frac{1}{(p^{d+1}; p)_{\ell+k}} \right] \frac{1}{(1-p)p^d} \frac{(p^a; p)_k (q^b; q)_{\ell} (p^c; p)_k}{(p; p)_k (q; q)_{\ell}} x^{\ell} y^k \\
&= \sum_{\ell, k=0}^{\infty} \left[1 - \frac{1-p^d}{1-p^{d+\ell+k}} \right] \frac{1}{(1-p)p^d} \frac{(p^a; p)_k (q^b; q)_{\ell} (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_k (q; q)_{\ell}} x^{\ell} y^k \\
&= \sum_{\ell, k=0}^{\infty} \left[\frac{1-p^{d+\ell+k}-1+p^d}{1-p^{d+\ell+k}} \right] \frac{(p^a; p)_k (q^b; q)_{\ell} (p^c; p)_k}{(1-p)p^d (p^d; p)_{\ell+k} (p; p)_k (q; q)_{\ell}} x^{\ell} y^k, \\
&= \frac{1}{1-p^d} \sum_{\ell, k=0}^{\infty} \frac{1-p^{\ell+k}}{1-p} \frac{(p^a; p)_k (q^b; q)_{\ell} (p^c; p)_k}{(p^c; p)_{\ell} (q^{d+1}; q)_k (p; p)_k (q; q)_{\ell}} x^{\ell} y^k \\
&= \frac{1}{1-p^d} \sum_{\ell, k=0}^{\infty} \left[\frac{1-p^k}{1-p} + \frac{p^k-p^{\ell+k}}{1-p} \right] \frac{(p^a; p)_k (q^b; q)_{\ell} (p^c; p)_k}{(p^c; p)_{\ell} (q^{d+1}; q)_k (p; p)_k (q; q)_{\ell}} x^{\ell} y^k \\
&= \frac{1}{1-p^d} \left[y D_{y,p}\Xi_1(p^{d+1}) + \frac{1}{1-p}\Xi_1(q^{d+1}, py) - \frac{1}{1-p}\Xi_1(q^{d+1}, px, py) \right].
\end{aligned}$$

Proceeding along the same lines as above, formulas (28)-(29) can be proved. $\square$

Theorem 2.7. For $0 < \Re(\alpha) < \Re(\gamma)$, the q -integral representations for Ξ_1 is true

$$\begin{aligned}
&\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\
&= \frac{1}{\Gamma_p(c)} \int_0^{\frac{1}{1-p}} E_p(-pu) u^{c-1} \Xi_2(p^a, q^b; p^d; p, q, x, (1-p)yu) d_p u.
\end{aligned} \tag{30}$$

Proof. Using the p -shifted factorials $(p^c; p)_k$ and the p -Gamma functions are defined as follows

$$(p^c; p)_k = \frac{(1-p)^k \Gamma_p(c+k)}{\Gamma_p(c)}$$

and

$$\Gamma_p(c) = \int_0^{\frac{1}{1-p}} E_p(-pu) u^{c-1} d_p u,$$

we obtain (30). $\square$

Theorem 2.8. For $s \in \mathbb{N}$, the differentiation formulas for Ξ_1 and Ξ_2 hold true

$$\mathbb{D}_{y,p}^s \left[y^{a+s-1} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \right] = \frac{(p^a; p)_s}{(1-p)^s} y^{a-1} \Xi_1(p^{a+s}, q^b, p^c; p^d; q, p, x, y), \tag{31}$$

$$\begin{aligned}\mathbb{D}_{x,q}^s \left[x^{b+s-1} \Xi_1(p^a, q^b, p^c; p^d; q, p, x, y) \right] &= \frac{(q^b; q)_s}{(1-q)^s} x^{b-1} \Xi_1(q^a, q^{b+s}, p^c; p^d; q, p, x, y), \\ \mathbb{D}_{y,p}^s \left[x^{c+s-1} \Xi_1(p^a, q^b, p^c; p^d; q, p, x, y) \right] &= \frac{(p^c; p)_s}{(1-p)^s} y^{c-1} \Xi_1(p^a, q^b, p^{c+s}; p^d; q, p, x, y)\end{aligned}\quad (32)$$

and

$$\begin{aligned}\mathbb{D}_{y,p}^s \left[y^{a+s-1} \Xi_2(p^a, q^b; p^c; p, q, x, y) \right] &= \frac{(p^a; p)_s}{(1-p)^s} y^{a-1} \Xi_2(p^{a+s}, q^b; p^c; q, p, x, y), \\ \mathbb{D}_{x,q}^s \left[x^{b+s-1} \Xi_2(p^a, q^b; p^c; q, p, x, y) \right] &= \frac{(q^b; q)_s}{(1-q)^s} x^{b-1} \Xi_2(q^a, q^{b+s}; p^c; q, p, x, y).\end{aligned}\quad (33)$$

Proof. Using

$$\mathbb{D}_{y,p}^s \left[y^{a+k+s-1} \right] = \frac{(p^{a+k}; p)_s}{(1-p)^s} y^{a+k-1},$$

and

$$(p^a; p)_k (p^{a+k}; p)_s = (p^a; p)_{k+s} = (p^a; p)_s (p^{a+s}; p)_k,$$

we get

$$\begin{aligned}&\mathbb{D}_{y,p}^s \left[y^{a+s-1} \Xi_1(p^a, q^b, p^c; p^d; q, p, x, y) \right] \\&= \sum_{\ell, k=0}^{\infty} \frac{(p^a; p)_k (q^b; q)_\ell (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^\ell \mathbb{D}_{y,p}^s \left[y^{a+k+s-1} \right] \\&= \sum_{\ell, k=0}^{\infty} \frac{(p^a; p)_k (q^b; q)_\ell (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^\ell \frac{(p^{a+k}; p)_s}{(1-p)^s} y^{a+k-1} \\&= \frac{1}{(1-p)^s} y^{a-1} \sum_{\ell, k=0}^{\infty} \frac{(q^b; q)_\ell (p^a; p)_{k+s} (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^\ell y^k \\&= \frac{(p^a; p)_s}{(1-p)^s} y^{a-1} \sum_{\ell, k=0}^{\infty} \frac{(p^{a+s}; p)_k (q^b; q)_\ell (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^\ell y^k \\&= \frac{(p^a; p)_s}{(1-p)^s} y^{a-1} \Xi_1(p^{a+s}, q^b, p^c; p^d; q, p, x, y).\end{aligned}$$

The proof Eqs. (32)-(33) are on the same lines as of Eq. (31). $\square$

Theorem 2.9. The summation formula for Ξ_1 and Ξ_2 holds true:

$$\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) = \sum_{\ell=0}^{\infty} \frac{(q^b; q)_\ell}{(p^d; p)_\ell (q; q)_\ell} x^\ell {}_2\phi_1(p^a, p^c; p^{d+\ell}; p, y) \quad (34)$$

and

$$\Xi_2(p^a, q^b; p^c; p, q, x, y) = \sum_{\ell=0}^{\infty} \frac{(q^b; q)_\ell}{(p^c; p)_\ell (q; q)_\ell} x^\ell {}_2\phi_1(p^a, 0; p^{c+\ell}; p, y). \quad (35)$$

Proof. From the definition of Ξ_1 and (3), we have

$$\begin{aligned}&\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\&= \sum_{\ell, k=0}^{\infty} \frac{(p^a; p)_k (q^b; q)_\ell (p^c; p)_k}{(p^d; p)_{\ell+k} (p; p)_k (q; q)_\ell} x^\ell y^k\end{aligned}$$

$$\begin{aligned}
&= \sum_{\ell=0}^{\infty} \frac{(q^b; q)_{\ell}}{(p^d; p)_{\ell}(q; q)_{\ell}} x^{\ell} \sum_{k=0}^{\infty} \frac{(p^a; p)_k (p^c; p)_k}{(p^{d+\ell}; p)_k (p; p)_k} y^k \\
&= \sum_{\ell=0}^{\infty} \frac{(q^b; q)_{\ell}}{(p^d; p)_{\ell}(q; q)_{\ell}} x^{\ell} {}_2\phi_1(p^a, p^c; p^{d+\ell}; p, y).
\end{aligned}$$

□

Theorem 2.10. *The summation formulas for Ξ_1 and Ξ_1 holds true:*

$$\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) = \frac{(p^a; p)_{\infty}}{(p^d; p)_{\infty}} \sum_{\ell, k=0}^{\infty} \frac{(q^b; q)_{\ell} (p^c; p)_k (p^{d+k}; p)_{\infty}}{(p^{d+k}; p)_{\ell} (p^{a+k}; p)_{\infty} (p; p)_k (q; q)_{\ell}} x^{\ell} y^k, \quad (36)$$

$$\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) = \frac{(p^c; p)_{\infty}}{(p^d; p)_{\infty}} \sum_{\ell, k=0}^{\infty} \frac{(p^a; p)_k (q^b; q)_{\ell} (p^{d+k}; p)_{\infty}}{(p^{d+k}; p)_{\ell} (p^{c+k}; p)_{\infty} (p; p)_k (q; q)_{\ell}} x^{\ell} y^k \quad (37)$$

and

$$\Xi_2(p^a, q^b; p^c; p, q, x, y) = \frac{(p^a; p)_{\infty}}{(p^c; p)_{\infty}} \sum_{\ell, k=0}^{\infty} \frac{(q^b; q)_{\ell} (p^{c+k}; p)_{\infty}}{(p^{c+k}; p)_{\ell} (p^{a+k}; p)_{\infty} (p; p)_k (q; q)_{\ell}} x^{\ell} y^k. \quad (38)$$

Proof. Using the identities

$$\begin{aligned}
(p^a; p)_k &= \frac{(p^a; p)_{\infty}}{(p^{a+k}; p)_{\infty}}, \\
(p^c; p)_k &= \frac{(p^c; p)_{\infty}}{(p^{c+k}; p)_{\infty}}, \\
(p^d; p)_k &= \frac{(p^d; p)_{\infty}}{(p^{d+k}; p)_{\infty}},
\end{aligned}$$

applying the q -binomial theorem

$$\begin{aligned}
\frac{(p^{d+k}; p)_{\infty}}{(p^{a+k}; p)_{\infty}} &= \sum_{r=0}^{\infty} \frac{(p^{d-a}; p)_s}{(p; p)_r} \binom{p^{a+k}}{p^r}, \\
\frac{(p^{d+k}; p)_{\infty}}{(p^{c+k}; p)_{\infty}} &= \sum_{r=0}^{\infty} \frac{(p^{d-c}; p)_s}{(p; p)_r} \binom{p^{c+k}}{p^r},
\end{aligned}$$

the familiar (Gaussian polynomials) q -binomial coefficient defined by

$${}_1\Phi_0(p^a; -; p, x) = \sum_{n=0}^{\infty} \frac{(p^a; p)_n}{(p; p)_n} x^n = \frac{(p^a x; p)_{\infty}}{(x; p)_{\infty}}; |x| < 1$$

and

$${}_0\Phi_0(-; -; p, y) = \sum_{k=0}^{\infty} \frac{1}{(p; p)_k} y^k = \frac{1}{(y; p)_{\infty}}, |y| < 1.$$

Substituting the above equations, we get

$$\begin{aligned}
&\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\
&= \sum_{\ell, k=0}^{\infty} \frac{(p^a; p)_k (q^b; q)_{\ell} (p^c; p)_k}{(p^d; p)_k (p^{d+k}; p)_{\ell} (p; p)_k (q; q)_{\ell}} x^{\ell} y^k \\
&= \sum_{\ell, k=0}^{\infty} \frac{(q^b; q)_{\ell} (p^c; p)_k (p^a; p)_{\infty} (p^{d+k}; p)_{\infty}}{(p^{d+k}; p)_{\ell} (p^d; p)_{\infty} (p^{a+k}; p)_{\infty} (p; p)_k (q; q)_{\ell}} x^{\ell} y^k
\end{aligned}$$

$$\begin{aligned}
&= \frac{(p^a; p)_\infty}{(p^d; p)_\infty} \sum_{\ell, k=0}^{\infty} \frac{(q^b; q)_\ell (p^c; p)_k (p^{d+k}; p)_\infty}{(p^{a+k}; p)_\infty (p^{d+k}; p)_\ell (p; p)_k (q; q)_\ell} x^\ell y^k \\
&= \frac{(p^a; p)_\infty}{(p^d; p)_\infty} \sum_{\ell, k, r=0}^{\infty} \frac{(q^b; q)_\ell (p^c; p)_k (p^{d-a}; p)_r}{(p^{d+k}; p)_\ell (p; p)_k (q; q)_\ell (p; p)_r} \left(p^{a+k}\right)^r x^\ell y^k \\
&= \frac{(p^a; p)_\infty}{(p^d; p)_\infty} \sum_{\ell, k=0}^{\infty} \frac{(q^b; q)_\ell (p^c; p)_k}{(p^{d+k}; p)_\ell (p; p)_k (q; q)_\ell} x^\ell y^k \sum_{r=0}^{\infty} \frac{(p^{d-a}; p)_r p^{(a+k)r}}{(p; p)_r} \\
&= \frac{(p^a; p)_\infty}{(p^d; p)_\infty} \sum_{\ell, k=0}^{\infty} \frac{(q^b; q)_\ell (p^c; p)_k}{(p^{d+k}; p)_\ell (p; p)_k (q; q)_\ell} x^\ell y^k {}_1\Phi_0(p^{d-a}; -, p, p^{a+k}).
\end{aligned}$$

Similarly, we can prove (37)-(38). $\square$

Theorem 2.11. *The following recursion formula holds true for Ξ_1*

$$\begin{aligned}
&\Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) \\
&= \sum_{\ell=0}^{\infty} \frac{(q^b; q)_\ell (yp^{a+c-d-\ell}; p)_\infty}{(p^d; p)_\ell (q; q)_\ell (y; p)_\infty} x^\ell {}_2\phi_1(p^{d+\ell-a}, p^{d+\ell-c}; p^{d+\ell}; p, yp^{a+c-d-\ell}) \\
&= \sum_{\ell=0}^{\infty} \frac{(q^b; q)_\ell (p^c; p)_\infty (yp^a; p)_\infty}{(p^d; p)_\ell (q; q)_\ell (p^{d+\ell}; p)_\infty (y; p)_\infty} x^\ell {}_2\phi_1(p^{d+\ell-c}, y; yp^a; p, p^c) \\
&= \sum_{\ell=0}^{\infty} \frac{(q^b; q)_\ell (p^{d+\ell-c}; p)_\infty (yp^c; p)_\infty}{(p^d; p)_\ell (q; q)_\ell (p^{d+\ell}; p)_\infty (y; p)_\infty} x^\ell {}_2\phi_1(yp^{a+c-d-\ell}, p^c; yp^c; p, p^{d+\ell-c}) \\
&= \sum_{\ell=0}^{\infty} \frac{(q^b; q)_\ell (yp^a; p)_\infty}{(p^d; p)_\ell (q; q)_\ell (y; p)_\infty} x^\ell {}_2\phi_2(p^a, p^{d+\ell-c}; p^{d+\ell}, yp^a; p, yp^c).
\end{aligned} \tag{39}$$

Proof. On the other hand, we shall need the Heine's transformations formulas for p -hypergeometric functions:(see)

$$\begin{aligned}
{}_2\phi_1(p^a, p^b; p^c; p, y) &= \frac{(yp^{a+b-c}; p)_\infty}{(y; p)_\infty} {}_2\phi_1(p^{c-a}, p^{c-b}; p^c; p, yp^{a+b-c}) \\
&= \frac{(p^b; p)_\infty (yp^a; p)_\infty}{(p^c; p)_\infty (x; p)_\infty} {}_2\phi_1(p^{c-b}, y; yp^a; p, p^b) \\
&= \frac{(p^{c-b}; p)_\infty (yp^b; p)_\infty}{(p^c; p)_\infty (y; p)_\infty} {}_2\phi_1(yp^{a+b-c}, p^b; yp^b; p, p^{c-b}) \\
&= \frac{(yp^a; p)_\infty}{(y; p)_\infty} {}_2\phi_2(p^a, p^{c-b}; p^c, yp^a; p, yp^b).
\end{aligned} \tag{40}$$

In order to derive a list of p -analogues of transformation formulas for bibasic Humbert hypergeometric functions. Replacing p^a , p^b and p^c in (40) by p^a , p^c and $p^{d+\ell}$, gives

$$\begin{aligned}
{}_2\phi_1(p^a, p^c; p^{d+\ell}; p, y) &= \frac{(yp^{a+c-d-\ell}; p)_\infty}{(y; p)_\infty} {}_2\phi_1(p^{d+\ell-a}, p^{d+\ell-c}; p^{d+\ell}; p, yp^{a+c-d-\ell}) \\
&= \frac{(p^c; p)_\infty (yp^a; p)_\infty}{(p^{d+\ell}; p)_\infty (y; p)_\infty} {}_2\phi_1(p^{d+\ell-c}, y; yp^a; p, p^c) \\
&= \frac{(p^{d+\ell-c}; p)_\infty (yp^c; p)_\infty}{(p^{d+\ell}; p)_\infty (y; p)_\infty} {}_2\phi_1(yp^{a+c-d-\ell}, p^c; yp^c; p, p^{d+\ell-c})
\end{aligned}$$

$$= \frac{(yp^a; p)_\infty}{(x; p)_\infty} {}_2\phi_2(p^a, p^{d+\ell-c}; p^{d+\ell}, yp^a; p, yp^c). \quad (41)$$

□

Remark 2.12. The basic functions Ξ_1 and Ξ_2 are a q -analogue of the Humbert confluent hypergeometric functions Ξ_1 and Ξ_2 defined by ([7], page 225, Equations (20)-(22)):

$$\lim_{q \rightarrow 1} \Xi_1(p^a, q^b, p^c; p^d; q, p, \frac{1-p}{1-q}x, \frac{1}{1-p}y) = \Xi_1(a, b; c, d; x, y) \quad (42)$$

and

$$\lim_{q \rightarrow 1} \Xi_2(p^a; q^b, q^c; q, p, \frac{1-p}{1-q}x, y) = \Xi_2(a, b; c, d; x, y). \quad (43)$$

Proof. Using the limit

$$\lim_{q \rightarrow 1} \frac{(q^\beta; q)_n}{(1-q)^n} = (\beta)_n,$$

we obtain (42)-(43). □

Theorem 2.13. The following relationship holds true between the basic functions Ξ_1 and Ξ_2 :

$$\lim_{c \rightarrow \infty} \Xi_1(p^a, q^b, p^c; p^d; p, q, x, y) = \Xi_2(p^a, q^b; p^d; q, p, x, y). \quad (44)$$

Proof. Using the limit

$$\lim_{c \rightarrow \infty} (p^c; q)_n = (0; q)_n = 1,$$

we obtain (44). □

§3 Some properties of new special cases

Here, we take $p = q$ in the above section and apply the techniques of the p -derivative operator for Humbert series Ξ_1 and Ξ_2 , simplification gives the desired results.

Theorem 3.1. The following equalities hold true:

$$[\Theta_{y,p}]_p \Xi_1 = \frac{[a]_p [c]_p}{[d]_p} y \Xi_1(p^{a+1}, p^{c+1}; p^{d+1}), \quad (45)$$

$$[\Theta_{x,p}]_p \Xi_1 = \frac{[b]_p}{(1-p)[d]_p} x \Xi_1(p^{b+1}; p^{d+1}),$$

$$[\Theta_{y,p}]_p \Xi_2 = \frac{[a]_p}{(1-p)[c]_p} y \Xi_2(p^{a+1}; p^{c+1}), \quad (46)$$

$$[\Theta_{x,p}]_p \Xi_2 = \frac{[b]_p}{(1-p)[c]_p} x \Xi_2(p^{b+1}; p^{c+1}).$$

Proof. Using the relations of (7),(8) and (9) and simplifying, we get (45) and (46). □

Theorem 3.2. The following equalities hold true:

$$\begin{aligned} [\Theta_{y,p} + a]_p \Xi_1 &= [a]_p \Xi_1(p^{a+1}), \\ [\Theta_{x,p} + b]_p \Xi_1 &= [b]_p \Xi_1(p^{b+1}), \\ [\Theta_{y,p} + c]_p \Xi_1 &= [c]_p \Xi_1(p^{c+1}), \\ [\Theta_{x,p} + \Theta_{y,p} + d - 1]_p \Xi_1 &= [d - 1]_p \Xi_1(p^{d-1}) \end{aligned} \quad (47)$$

and

$$\begin{aligned} [\Theta_{y,p} + a]_p \Xi_2 &= [a]_p \Xi_2 (p^{a+1}), \\ [\Theta_{x,p} + b]_p \Xi_2 &= [b]_p \Xi_2 (p^{b+1}), \\ [\Theta_{x,p} + \Theta_{y,p} + c - 1]_p \Xi_1 &= [c - 1]_p \Xi_2 (p^{c-1}). \end{aligned} \quad (48)$$

Proof. Using the relations (7) and (9), simplifying, we get (47)-(48). $\square$

Theorem 3.3. *The following equalities hold true:*

$$\left([\Theta_{y,p}]_p [\Theta_{x,p} + \Theta_{y,p} + d - 1]_p - y [\Theta_{y,p} + a]_p [\Theta_{y,p} + c]_p \right) \Xi_1 = 0, \quad (49)$$

$$\left([\Theta_{x,p}]_p [\Theta_{x,p} + \Theta_{y,p} + d - 1]_p - \frac{x}{1-p} [\Theta_{x,p} + b]_p \right) \Xi_1 = 0 \quad (50)$$

and

$$\begin{aligned} \left([\Theta_{y,p}]_p [\Theta_{x,p} + \Theta_{y,p} + c - 1]_p - \frac{y}{1-p} [\Theta_{y,p} + a]_p \right) \Xi_2 &= 0, \\ \left([\Theta_{x,p}]_p [\Theta_{x,p} + \Theta_{y,p} + c - 1]_p - \frac{x}{1-p} [\Theta_{x,p} + b]_p \right) \Xi_2 &= 0. \end{aligned} \quad (51)$$

Proof. Using the relations (7) and (9) and simplifying, we get (49)-(51). $\square$

Corollary 3.4. *The following relations hold true:*

$$\begin{aligned} & \left[\left(p^{d-1} p^{\Theta_{x,p}} - y p^{a+c} \right) [\Theta_{y,p}]_p [\Theta_{y,p}]_p + p^{d-1} [\Theta_{y,p}]_p [\Theta_{x,p}]_p \right. \\ & \quad \left. - \left(p^{d-1} - [d]_p + y(p^a [c]_p + p^c [a]_p) \right) [\Theta_{y,p}]_p - y[a]_p [c]_p \right] \Xi_1 = 0, \\ & \left[\left(p^{d-1} - y p^{a+c} \right) [\Theta_{y,p}]_p [\Theta_{y,p}]_p + p^{d-1} p^{\Theta_{y,p}} [\Theta_{y,p}]_p [\Theta_{x,p}]_p \right. \\ & \quad \left. - \left(p^{d-1} - [d]_p + y(p^a [c]_p + p^c [a]_p) \right) [\Theta_{y,p}]_p - y[a]_p [c]_p \right] \Xi_1 = 0, \\ & \left[p^{d-1} p^{\Theta_{x,p}} [\Theta_{x,p}]_p [\Theta_{y,p}]_p + p^{d-1} [\Theta_{x,p}]_p [\Theta_{x,p}]_p \right. \\ & \quad \left. - \left(p^{d-1} - [d]_p + \frac{x p^b}{1-p} \right) [\Theta_{y,p}]_p - \frac{x}{1-p} [b]_p \right] \Xi_1 = 0, \\ & p^{d-1} [\Theta_{y,p}]_p [\Theta_{x,p}]_p + p^{d-1} p^{\Theta_{y,p}} [\Theta_{x,p}]_p [\Theta_{x,p}]_p \\ & \quad - \left(p^{d-1} - [d]_p + \frac{x p^b}{1-p} \right) [\Theta_{x,p}]_p - \frac{x}{1-p} [b]_p \right] \Xi_1 = 0 \end{aligned} \quad (52)$$

and

$$\begin{aligned} & \left[p^{c-1} p^{\Theta_{x,p}} [\Theta_{y,p}]_p [\Theta_{y,p}]_p + p^{c-1} [\Theta_{y,p}]_p [\Theta_{x,p}]_p \right. \\ & \quad \left. - \left(p^{c-1} - [c]_p + \frac{y p^a}{1-p} \right) [\Theta_{y,p}]_p - \frac{y}{1-p} [a]_p \right] \Xi_2 = 0, \end{aligned}$$

$$\begin{aligned}
& p^{c-1} [\Theta_{y,p}]_p [\Theta_{y,p}]_p + p^{c-1} p^{\Theta_{y,p}} [\Theta_{y,p}]_p [\Theta_{x,p}]_p \\
& - \left(p^{c-1} - [c]_p + \frac{yp^a}{1-p} \right) [\Theta_{y,p}]_p - \frac{y}{1-p} [a]_p \Big] \Xi_2 = 0, \\
& \left[p^{c-1} p^{\Theta_{x,p}} [\Theta_{x,p}]_p [\Theta_{y,p}]_p + p^{c-1} [\Theta_{x,p}]_p [\Theta_{x,p}]_p \right. \\
& - \left. \left(p^{c-1} - [c]_p + \frac{xp^b}{1-p} \right) [\Theta_{y,p}]_p - \frac{x}{1-p} [b]_p \right] \Xi_2 = 0, \\
& p^{c-1} [\Theta_{y,p}]_p [\Theta_{x,p}]_p + p^{c-1} p^{\Theta_{y,p}} [\Theta_{x,p}]_p [\Theta_{x,p}]_p \\
& - \left(p^{c-1} - [c]_p + \frac{xp^b}{1-p} \right) [\Theta_{x,p}]_p - \frac{x}{1-p} [b]_p \Big] \Xi_2 = 0.
\end{aligned} \tag{53}$$

Proof. Using the relation

$$[\mu + \nu]_q = [\mu]_q + q^\mu [\nu]_q = [\nu]_q + q^\nu [\mu]_q \tag{54}$$

we obtain our relations (52) and (53). $\square$

Theorem 3.5. *The following recursion formulas hold true for Ξ_1 and Ξ_2*

$$\begin{aligned}
\Xi_1(p^a, p^b, p^c; p^d; p, x, y) &= \sum_{\ell=0}^{\infty} \frac{(p^b; p)_\ell}{(p^d; p)_\ell (p; p)_\ell} x^\ell {}_2\phi_1(p^a, p^c; p^{d+\ell}; p, y), \\
\Xi_1(p^a, p^b, p^c; p^d; p, x, y) &= \sum_{k=0}^{\infty} \frac{(p^a; p)_k (p^c; p)_k}{(p^d; p)_k (p; p)_k} y^k {}_2\phi_1(0, p^b; p^{d+k}; p, x)
\end{aligned} \tag{55}$$

and

$$\begin{aligned}
\Xi_2(p^a, p^b; p^c; p, x, y) &= \sum_{\ell=0}^{\infty} \frac{(p^b; p)_\ell}{(p^c; p)_\ell (p; p)_\ell} x^\ell {}_2\phi_1(p^a, 0; p^{c+\ell}; p, y), \\
\Xi_2(p^a, p^b; p^c; p, x, y) &= \sum_{k=0}^{\infty} \frac{(p^a; p)_k}{(p^c; p)_k (p; p)_k} y^k {}_2\phi_1(0, p^b; p^{c+k}; p, x).
\end{aligned} \tag{56}$$

Proof. Using relations (7) and (9) and simplifying, we get (55) and (56). $\square$

Corollary 3.6. *The following formulas hold true:*

$$\begin{aligned}
\Xi_1(p^a, p^b, p^c; p^d; p, x, 0) &= {}_2\phi_1(0, p^b; p^d; p, x), \\
\Xi_1(p^a, p^b, p^c; p^d; p, 0, y) &= {}_2\phi_1(p^a, p^c; p^d; p, y)
\end{aligned} \tag{57}$$

and

$$\begin{aligned}
\Xi_2(p^a, p^b; p^c; p, x, 0) &= {}_2\phi_1(0, p^b; p^d; p, x), \\
\Xi_2(p^a; p^b; p^c; p, 0, y) &= {}_2\phi_1(p^a, 0; p^c; p, y).
\end{aligned} \tag{58}$$

Proof. By using (6) and (7), and setting $x = 0$ and $y = 0$ in relations (55) respectively and simplifying we get (57), similarly by using (6) and (8), and setting $x =$ and $y = 0$ in relations (56) respectively and simplifying, we get the relation (58). $\square$

Acknowledgements

The authors are very grateful to the anonymous referees for many valuable comments and suggestions which helped to improve the paper.

Declarations

Conflict of interest The authors declare no conflict of interest.

References

- [1] G E Andrews. *Summations and transformations for basic Appell series*, J London Math Soc, 1972, 2(4): 618-622.
- [2] P Agarwal, S Jain, J Choi. *Certain q -series identities*, Rev Real Acad Cienc Exactas Fsicas Nat Ser A Mat, 2017, 111: 139-146.
- [3] W A Al-Salam, A Verma. *On quadratic transformations of basic series*, SIAM J Math Anal, 1984, 15: 414-420.
- [4] A Al édamat, A Shehata. *On Bibasic Humbert hypergeometric function Φ_1* , Malaysian Journal for Mathematical Sciences, 2023, 17(1): 77-86.
- [5] A Aral, V Gupta. *On the analogue of Stancu-Beta operators*, Applied Mathematics Letters, 2012, 25(1): 67-71.
- [6] A Aral, V Gupta, R P Agarwal. *Applications of q -Calculus in Operator Theory*, Springer, New York, 2013.
- [7] A Erdlyi, W Mangus, F Oberhettinger, F G Tricomi. *Higher Transcendental Functions-Vol. I*, McGraw-Hill Book Company, New York, Toronto and London, 1953.
- [8] T Ernst. *A Method for q -Calculus*, J Nonlinear Math Phys, 2003, 10(4): 487-525.
- [9] T Ernst. *A Comprehensive Treatment of q -Calculus*, Birkhauser, Basel, 2012.
- [10] S Z H Eweis, Z S I Mansour. *Generalized q -Bernoulli Polynomials Generated by Jackson q -Bessel Functions*, Results in Mathematics, 2022, 77(3): 1-37.
- [11] G Gasper. *Summation, Transformation, and Expansion formulas for bibasic series*, American mathematical society, 1989, 312(1): 257-277.
- [12] G Gasper, M Rahman. *Basic Hypergeometric Series*, Cambridge, Encyclopedia of Mathematics and its Applications, 2004.
- [13] G Gasper, M Rahman. *An Indefinite Bibasic Summation Formula and Some Quadratic, Cubic and Quartic Summation and Transformation Formulas*, Can J Math, 1990, XLII(1): 1-27.
- [14] H A Ghany. *q -Derivative of basic hypergeometric series with respect to parameters*, Int J Math Anal, 2009, 3(33): 1617-1632.
- [15] F H Jackson. *Basic double hypergeometric functions*, Quart J Math, 1944, 15: 49-61.
- [16] V K Jain. *Some expansions involving basic hypergeometric functions of two variables*, Pacific J Math, 1980, 91(2): 349-361.

- [17] M Mursaleen, S Tabassum, R Fatma. *On q -statistical summability method and its properties*, Iranian Jour Sci Tech Trans A: Sci, 2022, 46(2): 455-460.
- [18] N Mehmodov, V Gupta. *On certain q -analogue of Szász Kantorovich operators*, J Appl Math and Computing, 2011, 37: 407-419.
- [19] G V Milovanovi, V Gupta, N Malik. *(p, q) -Beta functions and applications in approximation*, Boletín de la Sociedad Matemática Mexicana, 2016, 24(1): 219-237.
- [20] K M Oraby, Z S Mansour. *On q -analogs of Struve functions*, Quaestiones Mathematicae, 2022, 18: 1-29.
- [21] S D Purohit. *Some recurrence relations for the generalized basic hypergeometric functions*, Bull Math Anal Appl, 2009, 1: 22-29.
- [22] M Rahman. *Some quadratic and cubic summation formulas for basic hypergeometric series*, Canad J Math, 1993, 45(2): 394-411.
- [23] V Sahai, A Verma. *Recursion formulas for q -hypergeometric and q -Appell series*, Commun Korean Math Soc, 2018, 33(1): 207-236.
- [24] V Sahai, A Verma. *n -th-order q -derivatives of Srivastavas general triple q -hypergeometric series with respect to parameters*, Kyungpook Mathematical Journal, 2016, 56(3): 911-925.
- [25] V Sahai, A Verma. *n th-Order q -derivatives of multivariable q -hypergeometric series with respect to parameters*, Asian-European Journal of Mathematics, 2014, 7(2): 1450019.
- [26] D B Sears. *On the transformation theory of basic hypergeometric functions*, Proc London Math Soc, 1951, s2-53(1): 158-180.
- [27] A Shehata. *On the (p, q) -Bessel functions from the view point of the generating functions method*, Journal of Interdisciplinary Mathematics, 2020, 23 (7): 1435-1448.
- [28] A Shehata. *On the (p, q) -Humbert functions from the view point of the generating functions method*, J Funct Spaces, 2020, Article ID: 4794571.
- [29] A Shehata. *On basic Horn hypergeometric functions H_3 and H_4* , Adv Difference Equ, 2020: 595, <https://doi.org/10.1186/s13662-020-03056-3>.
- [30] A Shehata. *On q -Horn hypergeometric functions H_6 and H_7* , Axioms, 2021, 10(4): 336, <https://doi.org/10.3390/axioms10040336>.
- [31] A Shehata. *Certain new formulas for bibasic Humbert hypergeometric functions Ψ_1 and Ψ_2* , arXiv:2207.01689.
- [32] A Shehata. *An extension of basic Humbert hypergeometric functions*, arXiv:2104.08407.
- [33] H M Srivastava, P W Karlsson. *Multiple Gaussian Hypergeometric Series (Ellis Horwood Series Mathematics and its Applications)*, Ellis Horwood Ltd, 1986, <https://doi.org/10.1112/blms/18.1.84>.
- [34] H M Srivastava, A Shehata. *A family of new q -Extensions of the Humbert functions*, European Journal of Mathematical Sciences, 2018, 4 (1): 13-26.

- [35] B Srivastava. *A note on certain bibasic q -Appell and Lauricella series*, Glas Mat, 1995, 30(50): 29-36.
- [36] B Srivastava. *Bibasic q -Appell Series Connected with Ramanujans continued fractions*, Tamkang J Math, 2007, 38(1): 51-56.
- [37] B Srivastava. *Some relations between contiguous bibasic q -Appell functions*, Notas de matematica Notas de Matematica, 1998, 173: 1-6.
- [38] B Srivastava. *On a two parameter family of generalised q -Exponential function*, Mathematical Journal of Okayama University, 2002, 44: 137-142.
- [39] A Verma, V Sahai. *Some recursion formulas for q -Lauricella series*, Afr Mat, 2020, 31: 643-686.
- [40] R K Yadav, S D Purohit, V K Vyas. *On transformations involving generalized basic hypergeometric function of two variables*, Rev Tc Ing Univ Zulia, 2010, 33(2): 176-182.

¹Provincial Key Laboratory of Data-Intensive Computing, Key Laboratory of Intelligent Computing and Information Processing, School of Mathematics and Computer Science, Quanzhou Normal University, Quanzhou 362000, China.

Email: qbcai@126.com

²Department of Mathematics, Al-Aqsa University, Gaza, Gaza Strip, Palestine.

Email: ghazikhamash@yahoo.com

³Department of Mathematics, Faculty of Science, Assiut University, Assiut 71516, Egypt.

Emails: shimaa1362011@yahoo.com, shimaa_m@science.aun.edu.eg

⁴Department of Mathematics, Unaizah College of Science and Arts, Unaizah 56264, Qassim University, Buraydah 52571, Qassim, Saudi Arabia.

Emails: drshehata2006@yahoo.com, aymanshehata@science.aun.edu.eg, A.Ahmed@qu.edu.sa