

Sweedler's duality of Hopf algebras in Yetter-Drinfeld's categories and an application

GU Yue¹ WANG Shuan-hong^{2,3,*}

Abstract. Let ${}^H_H\mathcal{YD}$ denote the Yetter-Drinfeld category over a Hopf algebra H with a bijective antipode. Let $(A, m, \mu, \Delta, \varepsilon, S)$ be a Hopf algebra in the category ${}^H_H\mathcal{YD}$. Then we mainly study Sweedler's duality A^0 of A and show that $(A^0, m^{op}, \varepsilon^*, \Delta^{op}, \mu^*, S^*)$ is again a Hopf algebra in ${}^H_H\mathcal{YD}$ generalizing the main result in [8]. As an application of our result, we show that the space of binary-linearly recursive sequences is closed under the quantum convolution product of binary-linearly recursive sequences.

§1 Introduction

The concept of a Hopf algebra contains a symmetry between its algebraic structure and its coalgebraic structure and has many important applications (see [1], [6], [7] and [12]). A theory of linearly recursive sequences related Hopf algebras was first studied in 1980 by Peterson and Taft (see [9]) and later investigated in some papers (see [8], [14], [15], [16], [18] and [20]).

Throughout, \mathbb{k} will denote a fixed field. All vector spaces will be over \mathbb{k} and all homomorphisms will be \mathbb{k} -linear maps. Let C be a coalgebra with a coproduct $\Delta : C \longrightarrow C \otimes C$. We will adopt the Sweedler's "sigma" notation about Δ :

$$\Delta(c) = \sum c_1 \otimes c_2$$

for all $c \in C$ (cf. [12]).

In what follows, let (A, m_A, μ_A) be an associative algebra over \mathbb{k} . Then we have the coalgebra A^0 over \mathbb{k} given by Sweedler in [12] as follows:

$$A^0 = \{ f \in A^* \mid \text{Ker } f \text{ contains a cofinite ideal, } \}$$

where A^* is the linear dual space, and a cofinite ideal is an ideal J in A possessing that A/J is finite-dimensional. Furthermore, when A is a Hopf algebra, A^0 has a natural Hopf algebra structure which was described in [12, Section 6.2]. However, generally, A^{00} is not isomorphic to A , i.e., the Pontryagin duality does not hold (see [17]).

Received: 2024-02-20. Revised: 2025-11-17.

MR Subject Classification: 16T05, 16W99.

Keywords: Hopf algebra, Yetter-Drinfeld category, braided Hopf algebra, binary-linearly recursive sequence.

Digital Object Identifier(DOI): <https://doi.org/10.1007/s11766-026-5164-x>.

Supported by the National Natural Science Foundation of China (12271089, 12471033), the Jiangsu Provincial Scientific Research Center of Applied Mathematics (BK20233002).

*Corresponding author.

For $f \in A^*$ and $a, b \in A$, one sets $(a \vdash f)(b) = f(ba)$ (similarly, $(f \dashv a)(b) = f(ab)$). Then one gets that A^* is a A - A -bimodule. The following states are equivalent: for any $f \in A^*$

- (1) $f \in A^0 = (m_A^*)^{-1}(A^* \otimes A^*)$;
- (2) $A \vdash f$ is a finite-dimensional vector space;
- (3) $\dim(f \dashv A) < \infty$;
- (4) $A \vdash f \dashv A$ is finite-dimensional.

We denote $\mathbf{LS}_{\mathbb{k}}$ the category of linear spaces and linear maps. Then $(\mathbf{LS}_{\mathbb{k}}, \otimes_{\mathbb{k}}, \mathbb{k})$ is a symmetric category. We can describe the Sweedler's duality in the following

Theorem A. A^0 forms a braided Hopf algebra in $\mathbf{LS}_{\mathbb{k}}$.

Given a Hopf algebra H with coproduct Δ , counit ϵ , and antipode S . Then one writes ${}_H\mathcal{M}$ the category of left H -modules. Similarly for ${}^H\mathcal{M}$ the category of left H -comodules.

For $V \in {}^H\mathcal{M}$, we write $\rho(v) = \sum v_{(-1)} \otimes v_0$ for all $v \in V$. A left-left Yetter-Drinfeld module M over H is an object both in ${}_H\mathcal{M}$ and ${}^H\mathcal{M}$ such that

$$\rho(h \cdot m) = \sum h_1 m_{(-1)} S h_3 \otimes h_2 \cdot m_0 \tag{1}$$

holds, for $m \in M$ and $h \in H$.

We have the category ${}^H_H\mathcal{YD}$ of left-left Yetter-Drinfeld modules. The category ${}^H_H\mathcal{YD}$ has the braiding given by

$$\tau_{M,N} : M \otimes N \longrightarrow N \otimes M, \quad m \otimes n \mapsto \sum m_{(-1)} \cdot n \otimes m_0. \tag{2}$$

We note that the category ${}^H_H\mathcal{YD}$ can provide a solution of quantum Yang-Baxter equation (cf. [6]). Some related further studies can be found in [4], [11] and [19].

In this paper, we will replace the category $\mathbf{LS}_{\mathbb{k}}$ in Theorem A by the category ${}^H_H\mathcal{YD}$, and then show the following main result.

Theorem B. Let A be a braided Hopf algebra in the category ${}^H_H\mathcal{YD}$. Then A^0 is a braided Hopf algebra in ${}^H_H\mathcal{YD}$.

This paper consists of four sections. In Section 2, we mainly show **Theorem B**, generalizing Ng and Taft's main result in [8].

In Section 3 and Section 4, we show that the space of binary-linearly recursive sequences is closed under the quantum convolution product of binary-linearly recursive sequences (see Theorem 4.1 and Theorem 4.5), generalizing the main result in [8] for the linear case.

In Section 5, one studies the structural properties of the space of binary linearly recursive sequences (see Theorem 5.4).

§2 The proof of Theorem B

In this section, H denotes a Hopf algebra with a bijective antipode S .

Given a bialgebra $(B, m_B, \mu_B, \Delta_B, \epsilon_B)$ in category ${}^H_H\mathcal{YD}$. Recall from [1] that B^* is a left H -module with

$$(h \cdot f)(b) = f(S(h) \cdot b)$$

with $h, b \in H$ and $f \in B^*$.

Similarly for the coaction of H on B^* given by

$$\rho(f)(b) = S^{-1}(b_{(-1)}) \otimes f(b_0),$$

with $h \in H, b \in B$ and $f \in B^*$.

Given two algebras A and B in ${}^H_H\mathcal{YD}$. One has the algebra $A \otimes B$ with the multiplication

$$(x \otimes y)(a \otimes b) = x(y_{(-1)} \cdot a) \otimes y_0 b$$

for $x, a \in A$ and $y, b \in B$.

The following two lemmas are easy results.

Lemma 2.1. With the above notation,

- (1) The \dashv is a left H -linear;
- (2) The \vdash is a left H^{cop} -linear.

Lemma 2.2. B^0 is an H -submodule of B^* .

Proposition 2.3. B^0 is a subalgebra of B^* .

Proof. Let $f, g \in B^*$. For $a, b \in B$, we compute

$$\begin{aligned} ((fg) \dashv a)(b) &= (fg)(ab) = (f \otimes g)\Delta(ab) \\ &= f(a_1(a_{2(-1)} \cdot b_1)g(a_{20}b_2)) \\ &= f(a_{2(-1)2} \cdot [(S^{-1}(a_{2(-1)1}) \cdot a_1)b_1])g(a_{20}b_2) \\ &= (S^{-1}(a_{2(-1)2}) \cdot f)[(S^{-1}(a_{2(-1)1}) \cdot a_1)b_1]g(a_{20}b_2) \\ &= [(S^{-1}(a_{2(-1)2}) \cdot f) \dashv (S^{-1}(a_{2(-1)1}) \cdot a_1)](b_1)(g \dashv a_{20})(b_2) \\ &= \Delta^*[(S^{-1}(a_{2(-1)}) \cdot (f \dashv a_1) \otimes g \dashv a_{20})](b), \end{aligned}$$

where we apply that B is an H -module algebra to the third equation above and use Lemma 2.1 to the final equation.

Therefore, we have

$$\begin{aligned} (fg) \dashv B &\subseteq \Delta^*[H \cdot (f \dashv B) \otimes g \dashv B] \\ &\subseteq \Delta^*[(H \cdot f) \dashv B] \otimes g \dashv B. \end{aligned}$$

By $f \in B^0$ and Lemma 2.2, one obtains $H \cdot f \in B^0$. Applying $f, g \in B^0$ and the right-hand side of the above containment is finite-dimensional, so $fg \in B^0$. Finally, it is straight to show that $\varepsilon_B^*(1) \in B^0$.

This concludes the proof. ■

Lemma 2.4. $i \circ \tau : (B^*)^{op} \otimes (B^*)^{op} \rightarrow (B \otimes B)^{*op}$ is a homomorphism as algebra in ${}^H_H\mathcal{YD}$

Proof. It follows the definition of the coaction of H on B^* . ■

Then we can get Theorem B as follows.

Theorem 2.5. Assume that $(B, m_B, \mu_B, \Delta_B, \varepsilon_B, S_B)$ is a Hopf algebra in ${}^H_H\mathcal{YD}$. Then $(B^0, (m_{B^0})^{op}, \varepsilon_B^*, (\Delta_{B^0})^{op}, \mu_B^*, S_B^*)$ is a braided Hopf algebra in ${}^H_H\mathcal{YD}$.

Proof. Following [8, Theorem 3.4], we have to finish checking the following steps:

- (1) B^0 is an H -subcomodule of B^* .
- (2) Observe that $(m_{B^0})^{op} = \Delta_B^* \circ i \circ \tau : B^0 \otimes B^0 \rightarrow B^*$. It is morphism in ${}^H_H\mathcal{YD}$. Obviously, $\varepsilon_B^* : \mathbb{k} \rightarrow B^0$ is. Thus, $(B^0, (m_{B^0})^{op}, \varepsilon_B^*)$ is an algebra in the category ${}_H\mathcal{M}$.
- (3) Note that $(\Delta_{B^0})^{op}$ is the composite map $B^0 \xrightarrow{m_B^*} i(B^0 \otimes B^0) \xrightarrow{(i \circ \tau)^{-1}} B^0 \otimes B^0$. This is morphism in ${}^H_H\mathcal{YD}$. $\mu_B^* : B^0 \rightarrow k$. So, $(B^0, (\Delta_{B^0})^{op}, \mu_B^*)$ is a coalgebra in the category ${}_H\mathcal{M}$.
- (4) $(\Delta_{B^0})^{op} : (B^0)^{op} \rightarrow (B^0)^{op} \otimes (B^0)^{op}$ is an algebra map.
- (5) $S_B^*(B^0) \subseteq B^0$.
- (6) $(m_{B^0})^{op}(S_B^* \otimes id_{B^0})(\Delta_{B^0})^{op} = \varepsilon_B^* \mu_B^*$ and $(m_{B^0})^{op}(id_{B^0} \otimes S_B^*)(\Delta_{B^0})^{op} = \varepsilon_B^* \mu_B^*$.

These checks are omitted here. ■

A Hopf algebra H is said to be quasitriangular provided there exists an invertible element $R = \sum R^{(1)} \otimes R^{(2)} \in H \otimes H$ satisfying ($r := R$),

$$(QT1) \sum R^{(1)}_1 \otimes R^{(1)}_2 \otimes R^{(2)} = \sum (R^{(1)} \otimes 1 \otimes R^{(2)})(1 \otimes r^{(1)} \otimes r^{(2)});$$

$$(QT2) \sum R^{(1)} \otimes R^{(2)}_1 \otimes R^{(2)}_2 = \sum (R^{(1)} \otimes 1 \otimes R^{(2)})(r^{(1)} \otimes r^{(2)} \otimes 1);$$

$$(QT3) \sum \Delta^{cop}(h) = R\Delta(h)R^{-1}, \text{ with } h \in H.$$

Furthermore, when $R^{-1} = \sum R^{(2)} \otimes R^{(1)}$, (H, R) is said to be triangular.

Dually, a coquasitriangular Hopf algebra is a pair $(H, \langle | \rangle)$, where H is a Hopf algebra over \mathbb{k} and $\langle | \rangle: B \otimes B \rightarrow \mathbb{k}$ is a k -linear form (braiding) which is convolution invertible in $Hom_{\mathbb{k}}(H \otimes H, \mathbb{k})$ such that the following compatibility relations hold, for any $h, g, l \in H$

$$(BR1) \langle h | gl \rangle = \langle h_1 | g \rangle \langle h_2 | l \rangle;$$

$$(BR2) \langle hg | l \rangle = \langle g | l_1 \rangle \langle h | l_2 \rangle;$$

$$(BR3) \langle h_1 | g_1 \rangle \langle g_2 h_2 \rangle = \langle h_1 g_1 | h_2 | g_2 \rangle.$$

If $\langle h_1 | g_1 \rangle \langle g_2 | h_2 \rangle = \epsilon(g)\epsilon(h)$, then $(H, \langle | \rangle)$ is called a cotriangular Hopf algebra.

Example 2.6. (1) Let (H, R) be a quasitriangular Hopf algebra. Then the category ${}^H\mathcal{M}$ left H -modules is a braided monoidal subcategory of ${}^H_H\mathcal{YD}$ with the following braiding

$$\tau_{M,N} : M \otimes N \longrightarrow N \otimes M, \quad m \otimes n \mapsto R^{(2)} \cdot n \otimes R^{(1)} \cdot m \tag{3}$$

for all $M, N \in {}^H\mathcal{M}$, $m \in M$ and $n \in N$.

Let A and B be algebras in ${}^H\mathcal{M}$. Then we have the braided product $A\#B$ (called braided algebra) of A and B given via

$$(a\#x)(b\#y) = a(R^{(2)} \cdot b)\#(R^{(1)} \cdot x)y \tag{4}$$

for any $a, b \in A$ and $x, y \in B$.

(2) Let $(H, \langle | \rangle)$ be a coquasitriangular Hopf algebra. Then the category ${}^H\mathcal{M}$ left H -comodules is a braided monoidal subcategory of ${}^H_H\mathcal{YD}$ with the following braiding

$$\tau_{M,N} : M \otimes N \longrightarrow N \otimes M, \quad m \otimes n \mapsto \langle m_{(-1)} | n_{(-1)} \rangle n_0 \otimes m_0 \tag{5}$$

for all $M, N \in {}^H\mathcal{M}$, $m \in M$ and $n \in N$.

Let A and B be algebras in ${}^H\mathcal{M}$. Then we have the braided product $A\#B$ (called braided algebra) of A and B given via

$$(a\#x)(b\#y) = \langle x_{(-1)} | b_{(-1)} \rangle ab_0\#x_0y \tag{6}$$

for any $a, b \in A$ and $x, y \in B$.

As a straightforward result of Theorem 1.5, we have

Corollary 2.7. ([8, Theorem 3.4]) Given a quasitriangular Hopf algebra (H, R) with a bijective antipode S . If $(A, m_A, \mu_A, \Delta_A, \epsilon_A, S_A)$ is a braided Hopf algebra in ${}^H\mathcal{M}$, then $(A^0, (m_{A^0})^{op}, \epsilon_A^*, (\Delta_{A^0})^{op}, \mu_A^*, S_A^*)$ is also a braided Hopf algebra in ${}^H\mathcal{M}$.

Theorem 2.8. Given a coquasitriangular Hopf algebra $(H, |)$ with a bijective antipode S . When $(A, m_A, \mu_A, \Delta_A, \epsilon_A, S_A)$ is a braided Hopf algebra ${}^H\mathcal{M}$, $(A^0, (m_{A^0})^{op}, \epsilon_A^*, (\Delta_{A^0})^{op}, \mu_A^*, S_A^*)$ is also a braided Hopf algebra in ${}^H\mathcal{M}$.

Finally, as an application, let $A = \mathbb{k}[x, y]$ be the bialgebra with x group-like element and with y $(x, 1)$ -primitive. Consider a cyclic group $G = \langle g \rangle$ of order n . One has a Hopf algebra $H = \mathbb{k}[G]$ with h group-like for any $h \in G$. Moreover, H is quasitriangular with $R = (1/n) \sum_{i,j=0}^{n-1} p^{-ij} (g^i \otimes g^j)$, where p is a primitive n th root of unity in \mathbb{k} (see [6] or [8]). Thus ${}^H\mathcal{M}$ is a braided monoidal

subcategory of ${}^H_H\mathcal{YD}$. We can study the Sweedler's duality $A^0 = \mathbb{k}[x, y]^0$ in Section 3.

§3 Binary-linearly recursive sequences

Let $A = \mathbb{k}[x]$ be the polynomial algebra in one variable x . It has a bialgebra structure given by $\Delta(x) = x \otimes 1 + 1 \otimes x$ and $\varepsilon(x) = 0$. On one hand, we can identify an element f in the dual space A^* with the sequences $(f_n)_{n \geq 0} = (f_0, f_1, f_2, \dots)$, where $f_n = f(x^n)$ for $n \geq 0$. On the other hand, A has a dual coalgebra $A^0 = \{f \in A^* \mid f(J) = 0 \text{ for some cofinite ideal } J \text{ of } A, \text{ i.e., } A/J \text{ is finite dimensional}\}$. Since a cofinite ideal J of $A = \mathbb{k}[x]$ is just a nonzero ideal generated by a monic polynomial $h(x) = x^r - h_1x^{r-1} - \dots - h_r$, the condition $f(J) = 0$ means that $f_n = h_1f_{n-1} + \dots + h_rf_{n-r}$ for all $n > r$. This means that f is linearly recursive, satisfying the recursive relation $h(x)$. Thus the space of linearly recursive sequences has a Hopf algebra structure. See the references [3],[9],[13],[14] and [15] for details of this structure.

Let $q \neq 0$ in \mathbb{k} . In 1997, Ng and Taft [8] showed that the space of linearly recursive sequences is closed under the quantum convolution product $(f_n) * _q (g_n) = (h_n)$, where $h_n = \sum_{i=0}^n \binom{n}{i}_q f_i g_{n-i}$ when q is a root of unity.

We now consider the bialgebra $A = \mathbb{k}[x, y]$ with x group-like element and with y a $(x, 1)$ -primitive. $A = \mathbb{k}[x] \otimes \mathbb{k}[y]$ as an algebra, and thus $A^0 = \mathbb{k}[x]^0 \otimes \mathbb{k}[y]^0$ as a coalgebra. We identify each f in A^* as a binary-sequence $(f_{i,j})$ for $i, j > 0$, where $f_{i,j} = f(x^i y^j)$. A row of such a binary-sequence is a sequence $\{f_{i,p} \mid p \geq 0\}$ for a fixed $i \geq 0$, which we say is parallel to the y -axis, or a sequence $\{f_{p,j} \mid p \geq 0\}$ for a fixed $j \geq 0$, which we say is parallel to the x -axis.

Let f be in A^0 , $f(J) = 0$ for a cofinite ideal J of A . For each i, j , the powers of x (y) spans a finite-dimensional space in A/J , so there is a minimal monic $h_i(x)$ ($h_j(y)$) in $\mathbb{k}[x]$ such that each row of f parallel to the y (x)-axis satisfies $h_i(x)$ ($h_j(y)$). Thus J contains the cofinite elementary ideal Γ generated by $h_i(x)h_j(y)$.

Recall from [8] that given a $q \neq 0$ in \mathbb{k} and an integer $n > 0$, one has the Gaussian polynomial for $0 \leq i \leq n$

$$\binom{n}{i}_q = \frac{(n)_q!}{(i)_q!(n-i)_q!},$$

where $(n)_q = 1 + q + \dots + q^{n-1}$ and the q -factorial of n is given by $(0)!_q = 1$ and

$$(n)!_q = (1)_q(2)_q \cdots (n)_q = \frac{(q-1)(q^2-1) \cdots (q^n-1)}{(q-1)^n}$$

if $n > 0$. It is a polynomial in q with coefficients in \mathbb{Z} . If $q = 1$, then $\binom{n}{i}$ is the ordinary binomial coefficient.

Let x and y be variables subject to the quantum plane relation $yx = qxy$. Then for any $n > 0$ we have

$$(x + y)^n = \sum_{0 \leq i \leq n} \binom{n}{i} x^i y^{n-i}. \tag{7}$$

Let $q = 1$. Then we have that $\binom{n}{k}$ is the ordinary binomial coefficient.

Proposition 3.1. ([8, Lemma 5.2]) Let q be a primitive n th root of 1. For integers $a \geq b \geq 0$, write $a = a'n + r, b = b'n + s$ for $0 \leq r, s < n$. Then

$$\begin{pmatrix} a \\ b \end{pmatrix}_q = \begin{pmatrix} a' \\ b' \end{pmatrix} \begin{pmatrix} r \\ s \end{pmatrix}_q,$$

where $\begin{pmatrix} r \\ s \end{pmatrix}_q = 0$ if $r < s$.

We will let $A = \mathbb{k}_q[x, y]$ with $yx = qxy$, here $0 \neq q \in \mathbb{k}$. Then $f \in A^*$ is regarded as the binary sequences $(f_{m,n})_{m,n \geq 0} = (f_{0,0}, f_{0,1}, \dots, f_{0,n}, f_{1,0}, \dots, f_{1,n}, \dots, f_{m,0}, \dots, f_{m,n}, \dots)$ where

$$f_{m,n} = f(x^m y^n) = q^{-mn} f(y^n x^m) = q^{-mn} f_{n,m}$$

for all $m, n \geq 0$. We call them the q -binary sequences. A has a bialgebra structure with a group-like element x and with a $(x, 1)$ -primitive element y , i. e., we have a comultiplication Δ given by the following

$$\begin{aligned} \Delta(x) &= x \otimes x, & \Delta(y) &= x \otimes y + y \otimes 1, \\ \varepsilon(x) &= 1, & \varepsilon(y) &= 0, \end{aligned}$$

requiring Δ and ε to be algebra homomorphisms from A to $A \otimes A$ and A to \mathbb{k} , respectively. Thus one has

$$\Delta(x^m y^n) = \sum_{0 \leq k \leq n} \begin{pmatrix} n \\ k \end{pmatrix}_q x^{m+k} y^{n-k} \otimes x^m y^k \tag{8}$$

for any $m, n \geq 0$. Therefore, the quantum convolution product on A^* is given by $f_{m,n} *_q g_{m,n} = h_{m,n}$, where $h_{m,n} = \sum_{0 \leq k \leq n} \begin{pmatrix} n \\ k \end{pmatrix}_q f_{m+k, n-k} \otimes g_{m,k}$ for $m, n \geq 0$.

By a cofinite ideal J of $A = \mathbb{k}_q[x, y]$ we mean a nonzero ideal generated by a monic binary polynomial

$$\begin{aligned} h(x, y) &= x^r y^s - h_{1,0} x^{r-1} y^s - \dots - h_{r,0} y^s \\ &\quad - h_{0,1} x^r y^{s-1} - h_{1,1} x^{r-1} y^{s-1} - \dots - h_{r,1} y^{s-1} \\ &\quad - h_{0,2} x^r y^{s-2} - h_{1,2} x^{r-1} y^{s-2} - \dots - h_{r,2} y^{s-2} \\ &\quad \dots \quad \dots \quad \dots \quad \dots \\ &\quad - h_{0,s} x^r - h_{1,s} x^{r-1} - \dots - h_{r,s}. \end{aligned}$$

By the condition $f(J) = 0$, we have the following cases.

Case 1: If $f(x^{m-r} h(x, y) y^{n-s}) = 0$, then we have a binary linearly recursive sequence $f = (f_{m,n})_{m \geq r, n \geq s}$ satisfying the recursive relation $h(x, y)$, where

$$\begin{aligned} f_{m,n} &= h_{1,0} f_{m-1,n} + h_{2,0} f_{m-2,n} + \dots + h_{r,0} f_{m-r,n} \\ &\quad + h_{0,1} f_{m,n-1} + h_{1,1} f_{m-1,n-1} + \dots + h_{r,1} f_{m-r,n-1} \\ &\quad + h_{0,2} f_{m,n-2} + h_{1,2} f_{m-1,n-2} + \dots + h_{r,2} f_{m-r,n-2} \\ &\quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \\ &\quad + h_{0,s} f_{m,n-s} + h_{1,s} f_{m-1,n-s} + \dots + h_{r,s} f_{m-r,n-s}. \end{aligned}$$

Case 2: If $f(x^{m-r} y^{n-s} h(x, y)) = 0$, then we have a parameterized binary linearly recursive

sequence $f = (f_{m,n})_{m \geq r, n \geq s}$ satisfying the recursive relation $h(x, y)$, where

$$\begin{aligned}
 f_{m,n} = & q^{-(n-s)}h_{1,0}f_{m-1,n} + q^{-2(n-s)}h_{2,0}f_{m-2,n} + \cdots + q^{-r(n-s)}h_{r,0}f_{m-r,n} \\
 & + h_{0,1}f_{m,n-1} + q^{-(n-s)}h_{1,1}f_{m-1,n-1} + q^{-2(n-s)}h_{2,1}f_{m-2,n-1} \\
 & + \cdots + q^{-r(n-s)}h_{r,1}f_{m-r,n-1} \\
 & + h_{0,2}f_{m,n-2} + q^{-(n-s)}h_{1,2}f_{m-1,n-2} + q^{-2(n-s)}h_{2,2}f_{m-2,n-2} \\
 & + \cdots + q^{-r(n-s)}h_{r,2}f_{m-r,n-2} \\
 & \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \\
 & + h_{0,s}f_{m,n-s} + q^{-(n-s)}h_{1,s}f_{m-1,n-s} + q^{-2(n-s)}h_{2,s}f_{m-2,n-s} \\
 & + \cdots + q^{-r(n-s)}h_{r,s}f_{m-r,n-s}.
 \end{aligned}$$

Case 3: If $f(h(x, y)x^{m-r}y^{n-s}) = 0$, then we have a parameterized binary linearly recursive sequence $f = (f_{m,n})_{m \geq r, n \geq s}$ satisfying the recursive relation $h(x, y)$, where

$$\begin{aligned}
 f_{m,n} = & h_{1,0}f_{m-1,n} + h_{2,0}f_{m-2,n} + \cdots + h_{r,0}f_{m-r,n} \\
 & + q^{-(m-r)}h_{0,1}f_{m,n-1} + q^{-(m-r)}h_{1,1}f_{m-1,n-1} + q^{-(m-r)}h_{2,1}f_{m-2,n-1} \\
 & + \cdots + q^{-(m-r)}h_{r,1}f_{m-r,n-1} \\
 & + q^{-2(m-r)}h_{0,2}f_{m,n-2} + q^{-2(m-r)}h_{1,2}f_{m-1,n-2} + q^{-2(m-r)}h_{2,2}f_{m-2,n-2} \\
 & + \cdots + q^{-2(m-r)}h_{r,2}f_{m-r,n-2} \\
 & \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \\
 & + q^{-s(m-r)}h_{0,s}f_{m,n-s} + q^{-s(m-r)}h_{1,s}f_{m-1,n-s} + \cdots + q^{-s(m-r)}h_{r,s}f_{m-r,n-s}.
 \end{aligned}$$

Remarks: (1) Set $h'_{i,j} = q^{-i(n-s)}h_{i,j}$ in Case 1. We then have a new monic binary polynomial $h'(x, y)$ as follows:

$$\begin{aligned}
 h'(x, y) = & x^r y^s - q^{n-s}h'_{1,0}x^{r-1}y^s - \cdots - q^{r(n-s)}h'_{r,0}y^s \\
 & - h'_{0,1}x^r y^{s-1} - q^{n-s}h'_{1,1}x^{r-1}y^{s-1} - \cdots - q^{r(n-s)}h'_{r,1}y^{s-1} \\
 & - h'_{0,2}x^r y^{s-2} - q^{n-s}h'_{1,2}x^{r-1}y^{s-2} - \cdots - q^{r(n-s)}h'_{r,2}y^{s-2} \\
 & \dots \quad \dots \quad \dots \quad \dots \\
 & - h'_{0,s}x^r - q^{n-s}h'_{1,s}x^{r-1} - \cdots - q^{r(n-s)}h'_{r,s}.
 \end{aligned}$$

In this situation, we get a new binary linearly recursive sequence $f' = (f'_{m,n})_{m \geq r, n \geq s}$ satisfying the recursive relation $h'(x, y)$, where

$$\begin{aligned}
 f'_{m,n} = & h'_{1,0}f'_{m-1,n} + h'_{2,0}f'_{m-2,n} + \cdots + h'_{r,0}f'_{m-r,n} \\
 & + h'_{0,1}f'_{m,n-1} + h'_{1,1}f'_{m-1,n-1} + \cdots + h'_{r,1}f'_{m-r,n-1} \\
 & + h'_{0,2}f'_{m,n-2} + h'_{1,2}f'_{m-1,n-2} + \cdots + h'_{r,2}f'_{m-r,n-2} \\
 & \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \\
 & + h'_{0,s}f'_{m,n-s} + h'_{1,s}f'_{m-1,n-s} + \cdots + h'_{r,s}f'_{m-r,n-s},
 \end{aligned}$$

which satisfies the relation $h'(x, y)$.

(2) Similarly, set $h''_{i,j} = q^{-j(m-r)}h_{i,j}$ in Case 2. We then have a new monic binary polynomial $h''(x, y)$, hence we can obtain a new binary linearly recursive sequence $f'' = (f''_{m,n})_{m \geq r, n \geq s}$ satisfying the recursive relation $h''(x, y)$,

Example 3.2. Let $h(x, y) = (x - a)(y - b)$ for any $a, b \in k$. Then we have a binary linearly recursive sequence

$$f_{m,n} = bf_{m,n-1} + af_{m-1,n} - abf_{m-1,n-1}$$

satisfying $h(x, y)$, for all $m \geq 1$ and $n \geq 1$. If take $f_{0,0} = 1, f_{0,i} = 0$ and $f_{i,0} = 0$ for all $i \in \mathbb{N}$, then one gets a binary linearly recursive sequence $(1, 0, 0, \dots, 0, -ab, -ab^2, -ab^3, \dots, 0, -a^2b, -a^2b^2, -a^2b^3, \dots)$.

Remark: In the case of $q = 1$, the Case 1, the Case 2 and the Case 3 are the same.

§4 Quantum convolution for $q = 1$

Let $q = 1$. Then we have $f_{m,n} = f_{n,m}$ for any $m, n \geq 0$ and $\binom{n}{k}$ is the ordinary binomial coefficient. In this section we will study the binary linearly recursive sequence in Case 1 satisfying the recursive relation $h(x, y)$ as in Section 2.

In what follows, by a method similar to that in [8] we can consider $A = \mathbb{k}[x, y]$ in ${}_H\mathcal{M}$ (see the end of Section 1) via

$$g^i \cdot (x^k y^l) = p^{i(k+l)} x^k y^l, \quad \text{for all } x^k, y^l \in A, g^i \in H. \tag{9}$$

It is not hard to verify that A is an algebra in ${}_H\mathcal{M}$. By (2) and (5), we have

$$\begin{aligned} & \tau_{A,A}(x^k y^l \otimes x^s y^t) \\ &= \frac{1}{n} \sum_{i,j=0}^{n-1} p^{-ij} (g^j \cdot (x^s y^t) \otimes g^i \cdot (x^k y^l)) \\ &= \frac{1}{n} \sum_{i,j=0}^{n-1} p^{-ij} p^{j(s+t)} p^{i(k+l)} (x^s y^t \otimes x^k y^l) \\ &= \frac{1}{n} \sum_{i,j=0}^{n-1} p^{-ij} p^{jr} p^{iu} (x^s y^t \otimes x^k y^l) \\ &= \frac{1}{n} \sum_{i=0}^{n-1} p^{iu} \left(\sum_{j=0}^{n-1} p^{j(r-i)} \right) (x^s y^t \otimes x^k y^l), \end{aligned}$$

where we write $s + t = an + r, k + l = bn + u$ for some $0 \leq r, u < n$.

If $i \neq r$, then $\sum_{j=0}^{n-1} p^{j(r-i)} = 0$. Therefore, $\tau_{A,A}(x^k y^l \otimes x^s y^t) = p^{(s+t)(k+l)} (x^s y^t \otimes x^k y^l)$. Thus we have a braided algebra $A \# A$ with

$$(x^{k_1} y^{l_1} \# x^{s_1} y^{t_1})(x^{k_2} y^{l_2} \# x^{s_2} y^{t_2}) = p^{(s_1+t_1)(k_2+l_2)} (x^{k_1+k_2} y^{l_1+l_2} \# x^{s_1+s_2} y^{t_1+t_2}) \tag{10}$$

for all $x^{k_1} y^{l_1} \# x^{s_1} y^{t_1}, x^{k_2} y^{l_2} \# x^{s_2} y^{t_2} \in A \# A$.

Note that $q = 1$, so $xy = yx$. It follows from (6) that $(x \# y)(y \# 1) = p(xy \# y) = p(y \# 1)(x \# y)$. If we regard A as a bialgebra in the category ${}_H\mathcal{M}$, then by Majid's bosonization [3], this requires Δ to be an algebra morphism in ${}_H\mathcal{M}$. Note that $(x \# y)^l = p^{\frac{1}{2}l(l+1)} (x^l \# y^l)$, and so, by (1) and (6) we have

$$\Delta(x^m y^l) = \sum_{0 \leq k \leq l} \binom{l}{k}_p p^{\frac{1}{2}[(l-k)(l-k+1)+m(m+2l+1)]} x^{m+l-k} y^k \# x^m y^{l-k} \tag{11}$$

for $m, l \geq 0$.

Notice that x^m is not a group-like element since $\Delta(x^m) = p^{\frac{1}{2}m(m+1)}(x^m \# y^m)$, and y^n is $(x^n, 1)$ -primitive, i.e., $\Delta(y^n) = x^n \# y^n + y^n \# 1$ since $\binom{n}{k}_p = 0$ for $1 \leq k \leq n - 1$. The counit ε of A is given as usual by $\varepsilon(x^m) = 1, \varepsilon(y^k) = \delta_{0,k}$. It is easy to check that Δ and ε are morphisms in ${}_H\mathcal{M}$. But, A is not a Hopf algebra in ${}_H\mathcal{M}$ unless $x^2 = 1$ with $S(x) = x$ and $S(y) = -xy$.

In what follows, Theorem 3.1 shows that the space A^0 of binary linearly recursive sequences is a bialgebra in ${}_H\mathcal{M}$. The quantum convolution product in A^* is given $(f_{m,l}) *_p (g_{m,l}) = (h_{m,l})$, where

$$h_{m,l} = \sum_{0 \leq k \leq l} \binom{l}{k}_p p^{\frac{1}{2}[(l-k)(l-k+1)+m(m+2l+1)]} f_{m+l-k,k} g_{m,l-k}. \tag{12}$$

Thus, we have

Theorem 4.1. Let p be a root of unity in k . Then the binary linearly recursive sequences is closed under the quantum convolution product $(f_{m,l}) *_p (g_{m,l}) = (h_{m,l})$ where $h_{m,l}$ is given by (12).

Remarks: 1) If $x = 1$ in $A = k[x, y]$, then by (7) we have

$$\Delta(y^l) = \sum_{0 \leq k \leq l} \binom{l}{k}_p y^k \# y^{l-k}$$

for all $l \geq 0$. Hence, we can get the result in [8, Theorem 4.1].

2) If $y = 1$ in $A = k[x, y]$, then by (7) one has

$$\Delta(x^m) = p^{\frac{1}{2}m(m+1)}(x^m \# y^m).$$

In this case, we have

$$h_m = p^{\frac{1}{2}m(m+1)} f_{m+l-k} g_m. \tag{13}$$

Notice that the algebraic structure of linearly recursive sequences under Hadamard product was described in [3]. We call the product given by (9) a quantum Hadamard product. Then we have

Corollary 4.2. The linearly recursive sequences are closed under the quantum Hadamard product $(f_m) *_p (g_m) = (h_m)$, where h_m is given by (13).

Next, we give a direct proof of Theorem 4.1.

First, we have

Lemma 4.3. Let $f = (f_{m,l})$ be the binary linearly recursive sequence over k satisfying the relation $h(x, y)$. Let $\alpha_1, \dots, \alpha_k; \beta_1, \dots, \beta_u$ be the roots of $h(x, y)$ in \bar{k} , the algebraic closure of k . Then, for any $i, j \geq 0, s, t \geq 0$, the subsequence $f^{(i,j)(s,t)}$ given by

$$\begin{aligned} f^{(i,j)(s,t)} = & (f_{i,s}, f_{i,s+t}, f_{i,s+2t}, \dots \\ & f_{i+j,s}, f_{i+j,s+t}, f_{i+j,s+2t}, \dots \\ & f_{i+2j,s}, f_{i+2j,s+t}, f_{i+2j,s+2t}, \dots \\ & \dots \dots \dots \dots) \end{aligned}$$

is a binary linearly recursive sequence over \bar{k} satisfying the relation $h_{j,t}(x, y) = (x - \alpha_1^j) \cdots (x - \alpha_k^j)(y - \beta_1^t) \cdots (y - \beta_u^t)$.

Proof. It suffices to show $h_{j,t}(x, y) \rightarrow f^{(i,j)(s,t)} = 0$ in order to verify that $f^{(i,j)(s,t)}$ satisfies $h_{j,t}(x, y)$, where \rightarrow is defined by $(a \rightarrow f)(b) = f(ba)$ for $a, b \in A$, and $f \in A^*$, for A is a commutative algebra.

Note that $f^{(i,j)(s,t)}(x^v y^w) = f_{i+vj, s+wt} = f(x^{i+vj} y^{s+wt})$ for $v, w \geq 0$. So, for all $g(x, y) \in k[x, y]$, $f^{(i,j)(s,t)}(g(x, y)) = f(x^i y^s g(x^j y^t))$. Thus, we have

$$\begin{aligned} h_{j,t}(x, y) &\rightarrow f^{(i,j)(s,t)}(x^v y^w) \\ &= f^{(i,j)(s,t)}(x^v y^w h_{j,t}(x, y)) \\ &= f(x^i y^s x^{vj} y^{wt} h_{j,t}(x^j, y^t)) \\ &= f(x^{i+vj} y^{s+wt} h_{j,t}(x^j, y^t)). \end{aligned}$$

For $h_{j,t}(x^j, y^t) = I(x, y)h(x, y)$ for some $I(x, y) \in k[x, y]$, and $h(x, y) \rightarrow f = 0$, one has $f(x^{i+vj} y^{s+wt} h_{j,t}(x^j, y^t)) = (h(x, y) \rightarrow f)(x^{i+vj} y^{s+wt} I(x, y)) = 0$, concluding the proof. ■

Example 4.4. In Example 2.2, we have a subsequence

$$\begin{aligned} f^{(1,2)(3,4)} &= (f_{1,3}, f_{1,7}, f_{1,11}, \dots \\ &\quad f_{3,3}, f_{3,7}, f_{3,11}, \dots \\ &\quad f_{5,3}, f_{5,7}, f_{5,11}, \dots) \end{aligned}$$

satisfying $h_{2,4}(x, y) = (x - a^2)(y - b^4)$. Explicitly, one has a binary linearly recursive sequence $g_{m,n} = b^4 g_{m,n-1} + a^2 g_{m-1,n} - a^2 b^4 g_{m-1,n-1}$ satisfying $h_{2,4}(x, y)$, and $(g_{m,n})_{m \geq 1, n \geq 1}$ is the same as $f^{(1,2)(3,4)}$. For example, $g_{0,0} = f_{1,3} = -ab^3$, $g_{0,1} = f_{1,7} = -ab^7$, $g_{1,0} = f_{3,3} = -a^3 b^3$, and it follows from the formula $g_{m,n}$ that

$$\begin{aligned} g_{1,1} &= b^4 g_{1,0} + a^2 g_{0,1} - a^2 b^4 g_{0,0} \\ &= b^4(-a^3 b^3) + a^2(-ab^7) - a^2 b^4(-ab^3) \\ &= -a^3 b^7 - a^3 b^7 + a^3 b^7 = -a^3 b^7, \end{aligned}$$

which equals to $f_{3,7}$.

Theorem 4.5. Let p be a primitive n th root of unity. Let $f = (f_{m,l})$ and $g = (g_{m,l})$ be binary linearly recursive sequences, satisfying $h(x, y) = \prod_{i,j} (x - \alpha_i)^{m_i} (y - \beta_j)^{n_j}$ and $J(x, y) = \prod_{k,r} (x - \gamma_k)^{s_k} (y - \lambda_r)^{t_r}$, respectively. Then the quantum product $(f_{m,l}) *_p (g_{m,l}) = (h_{m,l})$ where $h_{m,l}$ is given by (8), is a binary linearly recursive sequence, satisfying $W(x^n, y^n)$; here $W(x, y) = \prod_{i,k;j,r} [x - (\alpha_i^n + \gamma_k^n)^{m_i+s_k-1}] [y - (\beta_j^n + \lambda_r^n)^{n_j+t_r-1}]$.

Proof. For $0 \leq a, b \leq n$ and $c, d \geq 0$, we have

$$\begin{aligned} &h^{(a,n)(b,n)}(x^c y^d) \\ &= h_{a+cn, b+dn} \\ &= \sum_{u=0}^{b+dn} \binom{b+dn}{u}_p p^{\frac{1}{2}[(b+dn-u)(b+dn-u+1)+(a+cn)(a+2b+(c+2d)n+1)]} \\ &\quad f_{a+b+(c+d)n-u, u} g_{a+cn, b+dn-u} \quad (\text{by (8)}) \\ &= \sum_{s=0}^b \sum_{j=0}^d \binom{b+dn}{u}_p p^{\frac{1}{2}\{[(b-s)+(d-j)n][(b-s)+(d-j)n+1]+(a+cn)[a+2b+(c+2d)n+1]\}} \\ &\quad f_{a+b-s+(c+d-j)n, s+jn} g_{a+cn, b-s+(d-j)n} \end{aligned}$$

$$\begin{aligned}
 &= \sum_{s=0}^b \sum_{j=0}^d \binom{d}{j} \binom{b}{s} p^{\frac{1}{2}\{(b-s)+(d-j)n\}[(b-s)+(d-j)n+1]+(a+cn)[a+2b+(c+2d)n+1]} \\
 &\quad f_{a+b-s+(c+d-j)n, s+jn} g_{a+cn, b-s+(d-j)n} \quad (\text{by Proposition 3.1}) \\
 &= \sum_{s=0}^b \binom{b}{s} p^{\frac{1}{2}[(b-s)(b-s+1)+a(a+2b+1)]} \\
 &\quad \left(\sum_{j=0}^d \binom{d}{j} \right) (f^{(a+b-s,n)}(s,n))_{(c+d-j,j)} (g^{(a,n)}(b-s,n))_{(c,d-j)} \\
 &= \sum_{s=0}^b \binom{b}{s} p^{\frac{1}{2}[(b-s)(b-s+1)+a(a+2b+1)]} \\
 &\quad \left(\sum_{i=0}^d \binom{d}{i} \right) (f^{(a+b-s,n)}(s,n))_{(c+i,d-i)} (g^{(a,n)}(b-s,n))_{(c,i)}.
 \end{aligned}$$

Therefore,

$$h^{(a,n)}(b,n) = \sum_{s=0}^b \binom{b}{s} p^{\frac{1}{2}[(b-s)(b-s+1)+a(a+2b+1)]} f^{(a+b-s,n)}(s,n) * g^{(a,n)}(b-s,n),$$

where $*$ is the usual convolution product. It follows from Lemma 4.3 that $f^{(a+b-s,n)}(s,n)$ and $g^{(a,n)}(b-s,n)$ are binary linearly recursive sequences, satisfying $h_{n,n}(x,y) = \prod_{i,j} (x - \alpha_i^n)^{m_i} (y - \beta_j^n)^{n_j}$ and $J_{n,n}(x,y) = \prod_{k,r} (x - \gamma_k^n)^{s_k} (y - \lambda_r^n)^{t_r}$, respectively. Hence Theorem 4.5 holds. In fact, h is the interlacing of the sequences $h^{(0,n)}(0,n), h^{(0,n)}(1,n), \dots, h^{(0,n)}(n-1,n), \dots, h^{(1,n)}(0,n), \dots, h^{(1,n)}(n-1,n), \dots, h^{(n-1,n)}(0,n), \dots, h^{(n-1,n)}(n-1,n)$.

This finishes the proof. ■

§5 Structural properties of binary linearly recursive sequence space

Let p be a primitive n th root of 1, \mathcal{BL} the space of binary linearly recursive sequences. Denote the quantum convolution product on \mathcal{BL} by $*_p$. Define a map $\Phi : \mathcal{BL} \rightarrow \mathcal{BL}$ by $\Phi(f) = f^{(0,n)}(0,n)$, where $f^{(0,n)}(0,n) = (f_{0,0}, f_{0,n}, f_{0,2n}, \dots, f_{n,0}, f_{n,n}, f_{n,2n}, \dots, f_{2n,0}, f_{2n,n}, f_{2n,2n}, \dots)$. Obviously, Φ is surjective.

Let $f, g \in \mathcal{BL}$, $h = f *_p g$. By the proof of Theorem 4.5, we get $h^{(0,n)}(0,n) = f^{(0,n)}(0,n) * g^{(0,n)}(0,n)$. Hence $\Phi(f *_p g) = \Phi(h) = h^{(0,n)}(0,n) = f^{(0,n)}(0,n) * g^{(0,n)}(0,n) = \Phi(f) * \Phi(g)$. Therefore, we have

Lemma 5.1. Φ is a surjective algebra map of $(\mathcal{BL}, *_p)$ onto $(\mathcal{BL}, *)$.

Let $Z^{(i,j)}$ be the sequence with 1 in the (i, j) th position (coordinate) and zero elsewhere. Then, by (8) one has

$$\begin{aligned}
 \text{Lemma 5.2. (i)} \quad & Z^{(i,j)} *_p Z^{(s,t)} = \binom{t+j}{j} p^{\frac{1}{2}[t(t+1)+s(i+t+2j+1)]} Z^{(s,t+j)}, \\
 \text{(ii)} \quad & Z^{(0,j)} *_p Z^{(0,t)} = \binom{t+j}{j} p^{\frac{1}{2}t(t+1)} Z^{(0,t+j)}, \\
 \text{(iii)} \quad & Z^{(i,0)} *_p Z^{(s,0)} = p^{\frac{1}{2}s(s+1)} Z^{(s,0)},
 \end{aligned}$$

- (iv) $Z^{(i,0)} *_p Z^{(0,t)} = p^{\frac{1}{2}t(t+1)} Z^{(0,t)}$,
- (v) $Z^{(0,j)} *_p Z^{(s,0)} = p^{\frac{1}{2}s(s+2j+1)} Z^{(s,j)}$.

Lemma 5.3. The kernel of Φ is a nilpotent ideal of $(\mathcal{BL}, *_p)$.

Proof. By the definition of Φ , an element f in $\text{Ker } \Phi$ is of the form $\sum_{i,j \not\equiv 0 \pmod n} \alpha_{i,j} Z^{(i,j)}$ (an infinite sum), so an element in $(\text{Ker } \Phi)^n$ is of the form

$$\sum_{i_1, \dots, i_n; j_1, \dots, j_n \not\equiv 0 \pmod n} \alpha_{i_1, \dots, i_n; j_1, \dots, j_n} Z^{(i_1, j_1)} *_p \dots *_p Z^{(i_n, j_n)}.$$

It suffices to verify that $Z^{(i_1, j_1)} *_p \dots *_p Z^{(i_n, j_n)} = 0$ if $i_1, \dots, i_n; j_1, \dots, j_n \not\equiv 0 \pmod n$. Set $i_u = a_u n + r_u, j_v = b_v n + s_v$ with $0 < r_u, s_v < n$ for $u, v = 1, \dots, n$. There are $u', v' \leq n-1$ such that $r_1 + \dots + r_{u'} < n, s_1 + \dots + s_{v'} < n$, but $r_1 + \dots + r_{u'} + r_{u'+1} \geq n$, and $s_1 + \dots + s_{v'} + s_{v'+1} \geq n$. So, by Lemma 5.2(i), for some β in k , we have

$$\begin{aligned} & Z^{(i_1, j_1)} *_p \dots *_p Z^{(i_{u'+1}, j_{v'+1})} \\ &= \beta Z^{(i_{u'}, j_1 + \dots + j_{v'})} *_p Z^{(i_{u'+1}, j_{v'+1})} \\ &= \alpha \binom{j_1 + \dots + j_{v'} + j_{v'+1}}{j_{v'+1}}_p Z^{(i_{u'+1}, j_1 + \dots + j_{v'} + j_{v'+1})}, \end{aligned}$$

where $\alpha = \beta p^{\frac{1}{2}[j_{v'+1}(j_{v'+1}+1) + i_{u'+1}(i_{u'} + 2j_1 + \dots + 2j_{v'} + j_{v'+1} + 1)]}$. Note that $j_1 + \dots + j_{v'} + j_{v'+1} = n(b_1 + \dots + b_{v'+1}) + (s_1 + \dots + s_{v'+1} - n) + n = n(b_1 + \dots + b_{v'+1} + 1) + (s_1 + \dots + s_{v'+1} - n)$, here $s_1 + \dots + s_{v'+1} - n < n$. By Proposition 3.1 and an argument similar to [8, Lemma 5.5], we have $\binom{j_1 + \dots + j_{v'} + j_{v'+1}}{j_{v'+1}}_p = 0$ and so $(\text{Ker } \Phi)^n = 0$.

This concludes the proof. ■

Theorem 5.4. $f \in \mathcal{BL}$ is $*_p$ -invertible if and only if $\Phi(f)$ is $*$ -invertible.

Proof. Similar to [8, Theorem 5.6]. It follows from Lemma 5.1 and Lemma 5.3 that Theorem 5.4 holds. ■

Acknowledgements

The authors are grateful to the referee for very careful reading and suggesting comments.

Declarations

Conflict of interest The authors declare no conflict of interest.

References

- [1] S Caenepeel, F Van Oystaeyen, Y Zhang. *Quantum Yang-Baxter module algebras*, *K-Theory*, 1994, 8(3): 231-255.
- [2] L Delvaux, A Van Daele. *Algebraic quantum hypergroups*, *Adv Math*, 2011, 226(2): 1134-1167.
- [3] R G Larson, E J Taft. *The algebraic structure of linearly recursive sequences under Hadamard product*, *Israel J Math*, 1990, 72: 118-132.

- [4] L Liu, S Wang, G Shi, et al. *Infinitesimal H-pseudobialgebra and associative pseudo-Yang-Baxter equation*, Comm Algebra, 2022, 50(3): 1071-1091.
- [5] S Mac Lane. *Categories for the Working Mathematician*, New York: Springer, 1971.
- [6] S Majid. *Foundations of Quantum Group Theory*, Cambridge: Cambridge University Press, 1998.
- [7] S Montgomery. *Hopf Algebras and Their Actions on Rings*, American Mathematical Society, 1993.
- [8] S H Ng, E J Taft. *Quantum convolution of linearly recursive sequences*, J Algebra, 1997, 198(1): 101-119.
- [9] B Peterson, E J Taft. *The Hopf algebra of linearly recursive sequences*, Aequationes Math, 1980, 20: 1-17.
- [10] D Radford. *The structure of a Hopf algebra with a projection*, J Algebra, 1985, 92(2): 322-347.
- [11] G Shi, S Wang. *Schur-Weyl quasi-duality and (co)triangular Hopf quasigroups*, J Math Phys, 2020, 61(5): 051701.
- [12] M E Sweedler. *Hopf Algebras*, New York: W A Benjamin, 1969.
- [13] M Smith. *The Pontrjagin duality theorem in linear spaces*, Ann Math, 1952, 56(2): 248-253.
- [14] E J Taft. *Hurwitz invertibility of linearly recursive sequences*, Congr Number, 1990, 73: 37-40.
- [15] E J Taft. *Algebraic aspects of linearly recursive sequences*, In: J Bergen, S Montgomery. *Advances in Hopf algebras*, Boca Raton: CRC Press, 1994.
- [16] E J Taft. *Hadamard invertibility of linearly recursive sequences in several variables*, Discrete Math, 1995, 139(1): 393-397.
- [17] A Van Daele, S H Wang. *Duality for weak multiplier Hopf algebras with sufficiently many integrals*, Adv Math, 2024, 458: 109971.
- [18] S H Wang. *Lie bialgebra structure of multivariate linearly recursive sequences*, Chinese Sci Bull, 1996, 41(4): 271-275.
- [19] T Zhang, Y Gu, S H Wang. *Hopf quasimodules and Yetter-Drinfeld modules over Hopf quasigroups*, Algebra Colloq, 2021, 28(2): 213-242.
- [20] T Zhang, S H Wang, D G Wang. *A new approach to braided monoidal categories*, J Math Phys, 2019, 60(1): 013510.

¹School of Physical and Mathematical Sciences, Nanjing Tech University, Nanjing 211816, China.

Email: 202110006988@njtech.edu.cn

²School of Mathematics, Southeast University, Nanjing 210096, China.

³Shing-Tung Yau Center of Southeast University, Southeast University, Nanjing 210096, China.

Email: shuanhwang@seu.edu.cn